

No. 105 tota


# Boston Medical Library Association,

19 BOYLSTON PLACE,

Received Ahril 18.1897.

By Gift of W. Bowditch. M. D.

Bowdel 



# AN INTRODUCTION

TO

# CLINICAL MEDICINE

SIX LECTURES ON

THE METHOD OF EXAMINING PATIENTS,
AND THE MEANS NECESSARY FOR ARRIVING AT
AN EXACT DIAGNOSIS.

BY

JOHN HUGHES BENNETT, M.D., F.R.S.E.,

PROFESSOR OF THE INSTITUTES OF MEDICINE AND OF CLINICAL MEDICINE
IN THE UNIVERSITY OF EDINBURGH.

THIRD EDITION,

WITH NUMEROUS WOODCUTS.

EDINBURGH:
ADAM AND CHARLES BLACK.

MDCCCLVII.



1109

EDINBURGH: PRINTED BY R. AND R CLARK.



## PREFACE.

The sale of two large editions of this little work flatters me with the conviction that my object in publishing it has, to a great extent, been attained. The present edition has been carefully revised, contains much new matter, and is illustrated by an increased number of woodcuts. It will, therefore, I trust be even more useful than its predecessors to those who are earnestly endeavouring, in the schools of medicine, to overcome the preliminary difficulties of a laborious profession.

JOHN HUGHES BENNETT.

Edinburgh, December 1856.

Digitized by the Internet Archive in 2011 with funding from Open Knowledge Commons and Harvard Medical School

# CONTENTS.

## LECTURE I.

								Page
METHOD OF	Examin	ING PAT	IENTS		•			9
Circulato	ry Syst	em				*	•	11
Respirato	ry Syst	em	,		4			11
Nervous	System							12
Digestive	System	ı.					,	12
Genito-U	rinary	System			٠			12
Integume	entary S	ystem						13
Antecede	nt Histo	ry		•				13
Post-morte:	m Exan	ninations						17
External	Appear	ances						19
Head								19
Spinal Co	olumn							19
Neck								19
Chest								19
Abdomer	١.			,				20
Blood								20
Microsco	pic Exa	mination		•				20
Position	of the V	arious In	nternal	Organ	ıs			23
Inspection					• *			25
PALPATION		•		,				27
MENGUE AMO	AT.							20

## LECTURE II.


			Page
Percussion	٠		34
Different Sounds produced by Percussion .			36
Sense of Resistance produced by Percussion			37
General Rules to be followed			38
Special Rules to be followed			42
Percussion of the Lungs			48
Heart			48
Liver			5
Spleen			58
Stomach and Intestines .			54
Kidneys			57
Bladder			58
TEOMIDE III			
LECTURE III.			
Auscultation	٠		60
General Rules to be followed		•	60
Special Rules for Auscultating the Lungs .	•		62
Sounds elicited by the Pulmonary Organs			68
Alterations of the Natural Sounds	•		68
New or Abnormal Sounds	٠		66
Special Rules for Auscultating the Heart .			69
Sounds elicited by the Circulatory Organs.			70
Modifications of the Healthy Sounds .			72
New or Abnormal Sounds			74
Auscultation of the Abdomen			78
Auscultation of the Large Vessels			76
General Rules diagnostic of Pulmonary Diseases			77
General Rules diagnostic of Cardiac Diseases			79
LECTURE IV.			
THE USE OF THE MICROSCOPE	٠		82
Description of the Microscope	•		86
Mensuration and Demonstration	•		97
How to observe with a Microscope .			101

#### CONTENTS.

## LECTURE V.

									Lage
P	RINCIPAL API	PLICATIO	ONS OF	THE 1	Aicroso	COPE TO	DIAGN	osis	105
	Saliva					•			106
	Milk .								109
	Blood.								112
	Pus .		•						117
	Sputum		•						118
	Vomited Mat	ters	•			•			124
	Fœces								126
	Uterine and	Vaginal	Discha	arges					128
	Mucus								131
	Dropsical Flu	iids							132
	Urine								135
	Cutaneous E	ruptions	and U	lcers					139
Saliva Milk Blood Pus Sputum Vomited Matters Fœces Uterine and Vaginal Discharges Mucus Dropsical Fluids Urine Cutaneous Eruptions and Ulcers USE OF CHEMICAL TESTS  LECTURE VI.  CLASSIFICATION AND DIAGNOSIS OF SKIN DISEASES Classification Diagnosis		152							
			LEC	TUR	E VI				
C	LASSIFICATIO	n and I	DIAGNO	SIS OF	SKIN I	Disease	s		158
	Classification				•				159
	Diagnosis								165
	Daminola								el level





#### LECTURE I.

#### METHOD OF EXAMINING PATIENTS.

It is absolutely necessary that an examination of patients at the bedside should be conducted with order, and according to a well-understood plan. I have observed that some students, on being called upon, in their turn, to interrogate a case, have felt great embarrassment, and have been unable to proceed. Others have put their questions, as it were at random, without any apparent object, and wandered from one system of the economy to another, in a vain search for a precise diagnosis, and a rational indication of cure. But continual practice, and the adoption of a certain method, will remove all difficulty. No doubt, the questioning a patient, to arrive at a knowledge of his condition, requires as much skill in the medical practitioner, as examining a witness does in counsel at the bar. They make it an especial study, and you must do so likewise. You should remember that, in proportion as this duty is performed well or ill, is the probability that your opinion of the case may be correct or incorrect; and that, not only will the reputation you hold among your colleagues greatly depend on your ability in this matter, but that the public itself will promptly give its confidence to him whose interrogations reveal sagacity and talent.

The method of examination differs greatly among practitioners, and must necessarily vary in particular cases. Men of experience gradually form a certain plan of their own, which enables them to arrive at their object more rapidly and securely than that adopted with, perhaps, an equally good result by another. In a clinical class, however, and in order that every one present may follow and understand what is going forward, the method adopted must be uniform. I hold it to be a matter of great importance, that every one standing round the bed should take an equal interest in what is proceeding, and this he cannot do unless he is fully aware of the manner and object of the examination. The plan which appears to me the best, and which we shall follow, is the one I learnt when myself a clinical student in the wards of Professor Rostan, of Paris. Its object is to arrive, as quickly as possible, at a knowledge of the existing condition of the patient, in a way that will insure the examiner that no important organ has been overlooked or escaped notice.

For this purpose, we ascertain, in the first instance, the organ principally affected, and the duration of the disease, by asking two questions, "Where do you feel pain?" and, "How long have you been ill?" Let us suppose that the patient feels pain in the cardiac region, we immediately proceed to examine the heart functionally and physically, and then the circulatory system generally. We next proceed to those organs which usually bear the nearest relation to the one principally affected—say, the respiratory system—and we then examine the lungs functionally and physically. We subsequently interrogate the nervous, digestive, genito-urinary, and integumentary systems. It is a matter of little importance in what order these are examined—the chief point is, not to neglect any of them. Lastly, we in-

quire into the past history of the case, when we shall have arrived at all the information necessary for the formation of a diagnosis.

The following is the arrangement of symptoms and circumstances demanding attention under each of the seven heads into which the examination is divided:—

I. CIRCULATORY SYSTEM.—Heart—Uneasiness or pain; its action and rhythm; situation where the apex beats; extent of dulness determined by percussion; its impulse; murmurs—if abnormal, their character, and the position and direction in which they are heard loudest. Arterial pulse—Number of beats in a minute; large or small, strong or feeble, hard or soft, equal or unequal, regular or irregular, intermittent, confused, imperceptible, etc. If an aneurismal swelling exist, its situation, pulsations, symptoms, extent, and sounds, must be carefully examined. Venous pulse—If perceptible, observe position, force, etc.

II. RESPIRATORY SYSTEM.—Nares—Discharges; sneezing. Larynx and Trachea-Voice, natural or altered in quality, hoarse, difficulty of speech, aphonia, etc.; if affected, observe condition of epiglottis, tonsils, and pharynx, by means of a spatula. Lungs—State of respiration; easy or difficult, quick or slow, equal or unequal, laboured, painful, spasmodic, dyspnæa, etc.; odour of breath. Expectoration, trifling or profuse, easy or difficult; its character, thin or inspissated, frothy, mucous, purulent or muco-purulent, rusty, bloody; microscopical examination. Hæmoptysis, colour, appearance and amount of blood discharged. Cough, rare or frequent, short or long, painful or not, moist or dry. External form of the chest, unusually rounded or flattened, symmetrical or not, etc. Movements-regular, equal, their amount, etc. Resonance, as determined by percussion, increased or diminished, dulness, cracked-pot sound, etc.

Sounds determined by auscultation, if abnormal, their character and position.

- III. Nervous System.—Brain—Intelligence—augmented, perverted, or diminished; cephalalgia; hallucinations; delirium, stupidity, monomania, idiocy; sleep, dreams, vertigo, stupor, coma. Spinal cord and nerves—Pain in back; general sensibility, increased, diminished, or absent; special sensibility—sight, hearing, smell, taste, touch, their increase, perversion, or diminution; spinal irritation, as determined by percussion; motion, natural or perverted, fatigue, pain on movement; trembling, convulsions, contractions, rigidity, paralysis.
- IV. DIGESTIVE SYSTEM.—Mouth—Lips, teeth, and gums; taste in the mouth. Tongue—Mode of protrusion, colour, furred, coated, fissured, condition of papille, moist or dry. Fauces, tonsils, pharynx, and æsophagus—Deglutition—if impeded, examine the pharynx with a spatula, the cervical glands, neck, etc.; regurgitation. Stomach—Appetite, thirst, epigastric uneasiness or pain, swelling, nausea, vomiting, character of matters vomited, flatulence, eructations. Abdomen—Its measurement and palpation; pain, distension or collapse, borborygmi, tumours, constipation, diarrhæa, character of dejections, hæmorrhoids. Liver—Size, as determined by percussion, pain, jaundice, results of palpation, etc. Spleen—Size, as determined by percussion. If enlarged, examine blood microscopically.
- V. Genito-Urinary System.—Uterus—Condition of menstrual discharge, amenorrhea, dysmenorrhea, menorrhagia, leucorrhea, etc. If pain, or much leucorrheal discharge, examine os uteri and vagina with speculum; uterine or ovarian tumours; pain in back; difficulty in walking, or in defæcation; functions of mammæ. Kidney—Lumbar

pain; micturition; quantity and quality of urine, colour, specific gravity; precipitates, as determined by the microscope, and by chemical tests; action of heat; nitric acid, etc.; action on test papers; stricture; discharges from urethra; spermatorrhœa; etc.

VI. Integumentary System.—General posture; external surface; expression of countenance; obesity; emaciation; colour; rough or smooth; dry or moist; perspiration; marks or cicatrices; eruptions (see diagnosis of skin diseases); temperature; morbid growths or swelling; anasarca; œdema; emphysema; etc.

VII. ANTECEDENT HISTORY.—Age; parentage; constitution; hereditary disposition; trade or profession; place of residence; mode of living as regards food and drink; habits; epidemics and endemics; contagion and infection; exposure to heat, cold, or moisture; irregularities in diet; excesses of any kind; fatigue; commencement and progress of the disease; date of rigor or seizure; mode of invasion; previous treatment; in female cases whether married or single—have had children and miscarriages, previous diseases, etc.

Such are the principal points to which your attention should be directed during the examination of a case. A little practice will soon impress them on your memory, and in this manner habit will insure you that no very important circumstance has been overlooked. At first, indeed, it may appear to you that such a minute examination is unnecessary; but we shall have abundant opportunities of proving that, whilst a little extra trouble never does harm, ignorance of a fact frequently leads to error. It is surprising, also, how rapidly one thoroughly conversant with the plan, is able to examine a patient so as to satisfy himself that all the organs

and functions have been carefully interrogated. Remember that the importance of particular symptoms is not known to the patient, and that, consequently, it is not in his power voluntarily to inform you of the necessary particulars. It is always your duty to discover them.

In carrying out the examination, the following hints may be attended to:—

- 1. It should never be forgotten that you are examining a fellow-creature, who possesses the same sensitiveness to pain, and the same feelings as you do, and that everything that can increase the one or wound the other should be most carefully avoided. Prudence, kindness, and delicacy, are especially enjoined upon those who treat the sick.
- 2. The questions should be precise, simple, and readily comprehended. When an individual has a limited intelligence, or is accustomed to a particular dialect, you will not arrive at your object by becoming impatient, or talking in a loud voice, but by putting your interrogations in a clear manner, and in language proportioned to the intelligence of the individual.
- 3. It is often necessary, after asking the first question, "Where do you feel pain?" to tell the patient to put his or her hand on the part. An Irish peasant applies the term "heart" to an indefinite region, extending over great part of the chest and abdomen; and a woman, in speaking of pain in the stomach, often means the lower part of the abdomen.
- 4. When pain is referred to any circumscribed part of the surface, the place should always be examined by palpation, and, if possible, seen. Rostan relates very instructive cases where the omission of one or the other of these rules has led to curious errors in diagnosis.
- 5. Although the question "How long have you been ill?" is sufficiently plain, it is often difficult to determine the period of commencement of many diseases. In acute inflam-

matory or febrile disorders, we generally count from the first rigor. In chronic affections, a lengthened cross-examination is frequently necessary to arrive at the truth.

- 6. A state of fever may be said to exist when we find the pulse accelerated, the skin hot, the tongue furred, unusual thirst, and headache. These symptoms are commonly preceded by a period of indisposition, varying in duration, and ushered in by a rigor or sensation of cold. Such a febrile state may be idiopathic, when the case is called one of fever, or symptomatic of some local disease, when the nature of the case is determined by the organ affected and lesion present.
- 7. In endeavouring to ascertain the cause of the disease, great tact and skill in examination are necessary. We must guard ourselves against the preconceived views of the patient on the one hand, and be alive to the possibility of imposition on the other. Sometimes, with all our endeavours, no appreciable cause can be discovered; and at others we find a variety of circumstances, any one of which would be sufficient to occasion the malady.
- 8. In forming our diagnosis—that is, in framing a theory deduced from the facts elicited by examination—we should be guided by all the circumstances of the case, and be very careful that these are fully known before we hazard an opinion. Even then it is not always possible to come to a satisfactory conclusion, and in such cases the diagnosis should be deferred until further observation has thrown new light upon the nature of the disease.
- 9. In recording a case, it is, for the most part, only necessary to put down, under each head, the symptoms or signs present. If any system be quite healthy, it should be said that it is normal. In many cases, however, it is necessary to state what are called negative symptoms. This demands great tact, and exhibits a high degree of medical information. For instance, an attack of epilepsy generally commences

with a cry or scream; but sometimes there is none—when this should be stated. Again, no expectoration is a rare negative symptom in pneumonia. Symptoms which are usually present in the disease, but are absent in the particular case, constitute negative symptoms.

10. All mention of size should be, according to its exact measurement, in feet and inches. Situation is often referred to certain regions, into which the surface has been arbitrarily divided, such as subscapular, cardiac, epigastric, etc., but it is always better to refer at once to anatomical parts, such as the clavicle, particular rib, nipple, umbilicus, angle of scapula, and so on. Extent should also be determined by proximity to well-known fixed points. All vague statements, such as large, great, small, little, etc., should be carefully avoided; and in recording cases dates and references should always be given in the day of the month or of the disease, and not in the day of the week.

11. In conversing on, or discussing, the circumstances of the case at the bed-side, we should always use technical language. Thus instead of saying, a man has a cavern at the top of the lung, we should speak of a vomica under the clavicle; instead of saying, a man has diseased heart, we should speak of cardiac hypertrophy, or of insufficiency of the mitral or aortic valves, etc.

Having formed a diagnosis, and prescribed for the patient, the further examination should be conducted at intervals, varying, as regards time, according to the gravity of the case. In addition to the changes which may occur in the signs and symptoms previously noticed, the effects of remedies should be carefully inquired into, and care taken to ascertain whether the medicine and diet ordered have been administered. If the case prove fatal, the symptoms ushering in death, and the manner in which it occurs should be especially observed. Whenever a record of the case is to be kept, I cannot too strongly impress upon you the importance of

noting these down in a book at the time, rather than trusting to the memory.\* For a long series of years the reports of cases, dictated aloud by the professor, and written down at the bed-side by the clerk, has formed a leading feature of the Edinburgh system of clinical instruction, and constitutes the only trustworthy method of drawing up cases with accuracy.

When a patient dies, the examination is not completed. The time has now arrived when an inspection of the dead body confirms or nullifies the diagnosis of the observer. You should consider this as a most important part of the clinical course. It is invariably regarded with the greatest interest by those who practise their profession with skill. It is only in this manner that any errors they may have committed can be corrected; that the value of physical diagnosis can be demonstrated and properly appreciated, and the true nature or pathology of diseases, and the mode of treating them rationally, can ever be discovered.

But here, again, method and order are as necessary in the examination of the dead as of the living, and it is of equal importance that no viscus be overlooked. The three great cavities should always be investigated. Nothing is more injurious to the scientific progress of medicine than the habit of inspecting only one of them, to satisfy the curiosity of the practitioner, or to determine his doubts on this or that point. Many medical men direct their attention to a certain class of diseases, and are apt to attribute too much importance to a particular lesion. It has frequently happened to me, when pathologist to this institution, to observe, that after the physician has examined this or that organ, to which he has attributed the death of his patient, and left the theatre,

<sup>\*</sup> I have arranged a note-book for taking cases, according to the system of examination here recommended, which may be procured of Mr. Thinn, medical bookseller, close to the Infirmary.

that further examination, according to the routine I always practised, has revealed important lesions that were never suspected. Thus, a person supposed to die of Bright's disease of the kidney, may have a pneumonia that was latent and overlooked. Large caverns and tubercular deposits in the lungs may satisfy the physician, and he may leave the body, when intense peritonitis may be subsequently found, arising from intestinal perforation. A man has hypertrophy, with valvular disease of the heart; he dies suddenly, and everything is referred to the cardiac lesion. On opening the head, an apoplectic extravasation or yellow softening may be discovered. I cannot too strongly, therefore, impress upon you the necessity of always making a thorough post-mortem examination, and for this purpose you should if possible, obtain permission to inspect the body, and not any particular cavity.

The object of a post-mortem examination is threefold, -1st, the cause of death; 2d, an appreciation of the signs and symptoms; 3d, the nature of the disease. These inquiries are very distinct, but practitioners generally have only in view the two first. It frequently happens that, on the discovery of a lesion that seems to explain the fatal termination, they feel satisfied, and there is an end to the investigation. In medico-legal cases, this is the only object. But even here it is necessary to examine all the organs, to avoid a possibility of error, for how can any conscientious man form an opinion, that an abdominal disease has been fatal, if he be not satisfied by inspection that the chest and brain are healthy? Again, it often occurs that a particular sign or symptom is unusual or mysterious, and this, if explained by the examination, is sufficient for the practitioner. But it must be obvious that this throws no light upon the nature of the disease, or its mode of cure. To do this, morbid changes must be sought for, not in that advanced stage where they cause death, or occasion prominent symptoms, but at the very earliest period that can be detected. Hence we must call in the microscope to our assistance, and with its aid follow the lesion into the ultimate tissue of organs; we must observe the circumstances which produced it, as well as the symptoms and physical signs to which it gives rise, the secondary disorders, and the order of their sequence; their duration and mode of termination. This is the kind of extended investigation which can alone be serviceable to the advancement of medicine, and such, I trust, will be the object all of you will have in view in examining dead bodies. At all events, such are the views that I shall constantly endeavour to place before you during this course of clinical instruction.

The following is an arrangement of the organs, textures, etc., which demand your attention:—

- I. EXTERNAL APPEARANCES.—Number of hours after death. General aspect and condition of the body; peculiarities of person; marks on the surface; sugillation; amount of decomposition. In cases of suspected death by violence, great minuteness in the external examination is necessary.
- II. HEAD.—Scalp; calvaria; meninges; sinuses; choroid plexus; brain, its form and weight; cerebellum, its weight; cortical and medullary substance of brain; ventricles, exact quantity of fluid in each, which should be removed with a pipette—its character; medulla oblongata; nerves, and arteries at the base of the brain; base of cranium; sinuses.
- III. Spinal Column.—Integuments over spine; Vertebræ; Meninges; Cord; Nerves.
- IV. Neck.—Thyroid gland; larynx and its appendages; trachæa; tongue; tonsils; pharynx, œsophagus.
- V. Chest.—Thymus gland; lining membrane of bronchi; bronchial glands; pleuræ; contents of pleural cavity;

parenchyma of lungs; large thoracic veins; pericardium, its contents; general aspect and position of the heart; its weight; amount of blood in its various cavities; right auricle; coronary veins; auricular septum; right ventricle, size of its cavity; thickness and degree of firmness of its walls; endocardium; tricuspid valve; pulmonary artery, its calibre; pulmonary veins; left auricle; mitral valve; left ventricle; thickness and condition of its muscular tissue; size of its cavity; sigmoid valves; coronary arteries; aortic opening and arch; large arteries of the neck; thoracic aorta, its structure and calibre.

VI. ABDOMEN.—Peritoneum and peritoneal cavity; omentum; position of abdominal viscera; stomach; duodenum; small and large intestines; liver, its weight, form, and structure—its artery, veins, and ducts; gall bladder and its contents; portal system; pancreas and its duct; mesenteric and other absorbent glands; spleen, its weight, size, and structure; supra-renal capsules; kidneys, weight of each; secreting and excreting portions; pelvis; ureters; bladder; with the prostate and urethra in the male; in the female, uterus, ovaries, Fallopian tubes, vagina; abdominal aorta and vena cava; large abdominal arteries and veins; ganglia of the sympathetic system.

VII. Bloop.—Appearance in the cavities of the heart, in aorta, vena cava, vena portæ, etc.; coagulated and fluid portions—adhesion or not of the former.

VIII.—MICROSCOPIC EXAMINATION of all the morbid structures and fluids, the blood, etc. etc.

In carrying out the post-mortem examination, the following hints may be attended to:—


1. As I have already said, the head, chest, and abdomen should always be examined, but the spinal cord and neck need not be disturbed unless the symptoms indicate some lesion there. In special cases, particularly judicial ones,

however, every part should be carefully inspected, and in them it may be further necessary to investigate a variety of circumstances connected with external or surgical lesion, such as fractures, wounds, and burns; injury to the large vessels; alterations of the organs of sense, etc. etc.

- 2. Great care should be taken never to disfigure the body. Incisions through the skin, therefore, should be made in such directions that when the edges are afterwards sewn together, the necessary dissections below may not be visible. Neither should the body be exposed more than is needful, and delicacy demands that the genitals should always be kept covered. The wishes and feelings of friends and relations should invariably be held in consideration.
- 3. Before removing the stomach, or any portion of the intestines, a ligature should be placed above and below the tube, which should afterwards be opened with the greatest care, and the character of the contents, whether gaseous, fluid, mucous, bloody, fœcal, or containing foreign substances, observed before washing and inspecting the mucous surface. This rule should be especially followed in all medicolegal investigations, in which, from neglect of it, the ends of justice have been frequently defeated.
- 4. You should seize every opportunity of opening dead bodies with your own hands, and acquiring dexterity in exposing the cavities, taking out the viscera, etc. Nothing is more painful than to see the brain cut into or contused, in removing the calvarium; or the large vessels at the root of the neck wounded in disarticulating the sternum, so that the surrounding parts are deluged with blood; or the cardiac valves cut through, instead of being simply exposed; or awkward incisions made into the intestines, whereby fœces escape; slipping of ligatures, etc. etc. Coolness, method, knowledge of anatomy, and skilfulness in dissection, are as necessary when operating on the dead as on the living body.

- 5. In examinations made at private houses, it is not always necessary to remove the viscera. The heart, lungs, liver, kidneys, etc., may be readily examined in situ. But in this Infirmary, where every facility exists, the viscera are invariably taken out, and after describing the morbid alterations they present, I shall always pass them round, so that every one present may examine them.
- 6. It is a good rule never to omit the examination of a morbid texture or product microscopically, until experience has made you perfectly familiar with its minute structure.
- 7. Notes of the examination should always be made at the time. If organs are healthy, this should be distinctly stated, so that hereafter all doubt as to their having been carefully examined may be removed. Here negative appearances are often of as much consequence as negative symptoms.
- 8. In describing morbid appearances, we should be careful to state the physical properties of an organ or texture, such as the size, form, weight, density, colour, position, etc.; and avoid all theoretical language, such as its being inflamed, tubercular, or cancerous, as well as such indefinite description as small and large, narrow and wide, increased or diminished, etc. etc.; size should always be stated in feet and inches, and the amount of fluid in quarts, pints, or ounces.
- 9. The amount of care and time bestowed on the examination of an individual body, will vary according to circumstances. In some cases it may require continued investigation, involving microscopical and chemical research for several days. I have never heard of a student regretting the employment of too much care in post-mortem investigation, although the occurrence of omissions from carelessness and unacquaintance with morbid anatomy, are unfortunately too often exhibited by medical men in courts of justice, to the detriment of our profession in the eyes of the

public, and not unfrequently to the perversion or suppression of justice.




For the correct examination of the patient in the manner described, it will be found necessary to possess an accurate knowledge of the relative position of the various internal organs. These, it should be remembered, do not occupy exactly the same position after death as in the living body. "Expiration is the last act of life, and this last expiration is usually more extensive and forced than the expiration of tranquil life. In the dead body, the lungs

Fig. 1. View of internal organs after removal of the thoracic and abdominal parietes. (After Sibson.)

shrink up within the position that they usually occupy during life; at the same time the heart and its vessels retract, and the abdominal organs follow the diaphragm somewhat upwards."—(Sibson.)

The remarkable changes which occasionally occur in the natural position of the internal viscera may be judged of from a case recorded in the Pathological Transactions of London, vol. iv., by Dr. Miller, in which the stomach was distended in the manner represented in the accompanying woodcut.



Besides the method of general examination previously detailed, it is further essential to employ various special modes of investigation. These are inspection, palpation, mensuration, percussion, auscultation, the use of the micro-

Fig. 2. Case of great enlargement of the stomach. (After Miller and Humby.)

scope, and of chemical tests. I propose giving you a short account of each of these modes of exploration.

#### INSPECTION.

Inspection of a part or of the whole surface of the body in various positions is often of the utmost importance. The latter is necessary in the examination of army recruits, but can seldom be carried out rigorously in hospital, and still less in private practice. Delicacy forbids it in females. The part affected, however, ought always to be seen, a neglect of which rule has led to numerous errors. The various eruptions which appear on the surface of the body are spoken of in Lecture vi. Here I shall only shortly allude to the inspection of the thorax, and of the abdomen.

Inspection of the Thorax.—This refers to the form and configuration of the entire thorax, or its various parts, and to a careful comparison of the two sides when at rest and when in motion. It is often difficult, in cases where changes are not well marked, to determine them by mere inspection. To do so, a good light, and a proper position, both of the observer and of the patient, are necessary. The observer should, if possible, be directly in front of the patient, and whenever the case admits of it, the latter should be in the sitting posture. The chest may be so altered in disease as to be irregular or unsymmetrical, from alterations, congenital or acquired, in the form of the bones of the vertebral column or thoracic walls. Various portions of it may be expanded or bulge out, as in cases of empyema or thoracic tumours; or it may be retracted and depressed, as occurs in chronic phthisis. A case presented itself to the late Dr. Spittal at the Royal Dispensary of this city, where the retraction was so great on one side, that the student in charge of it had placed compresses on the sound side, in the belief that the healthy prominence there was indicative of a tumour. The motions of the chest bear reference to inspiration and expiration,\* which pass imperceptibly into one another, and can be made more rapid or prolonged voluntarily. A forced inspiration gives rise to more thoracic movement in the female than in the male, in whom it is more abdominal. In disease these motions are altered in various ways—1st, By general excess or diminution, as in cases of spasmodic asthma or laryngeal obstruction. 2d, By partial immobility, as in pleurisy; or by augmented expansion, as in the side opposite a pneumonia or pleurisy. 3d, By increased rapidity, as in pericarditis; or unusual slowness, as in comatose diseases.

Inspection of the Abdomen.—The abdomen in health is slightly convex, and marked by elevations and depressions, corresponding to the muscles in its walls, the umbilicus, and prominences of the viscera below. It varies according to age and sex—in youth being smoother and flatter than in the adult, and in females being broader inferiorly than in males, from the greater width of the pelvis. In disease

<sup>\* &</sup>quot;During inspiration, the clavicles, first ribs, and through them the sternum and all the annexed ribs, are raised; the upper ribs converge, the lower diverge; the upper cartilages form a right angle with the sternum, and the lower cartilages of opposite sides, from the seventh downwards, move further asunder, so as to widen the abdominal space between them, just below the ziphoid cartilage: the effect being to raise, widen, and deepen the whole chest, to shorten the neck, and apparently to lengthen the abdomen. During expiration, the position of the ribs and cartilages is reversed; the sternum and ribs descend; the upper ribs diverge, the lower converge; the upper cartilages form a more obtuse angle with the sternum, and the lower cartilages of opposite sides approximate, so as to narrow the abdominal space between them, just below the ziphoid: the effect being to lower, narrow, and flatten the whole chest, to lengthen the neck, and apparently to shorten the abdomen. It is to be observed, that during inspiration, while the ribs and sternum are moving upwards, the lungs and heart, and the abdominal organs, are moving downwards, and that, consequently, viewed in relation to the ribs, the descent of the internal organs appears to be greater than it really is."—(Sibson).

it may be, 1st, enlarged generally and symmetrically, as in dropsies, or partially and irregularly, from ovarian, hepatic, splenic, and other tumours; 2d, it may be retracted—generally from extreme emaciation, or partially, from local intestinal obstruction. The superficial abdominal veins are sometimes greatly enlarged, and at others distinct pulsations are visible, dependent on deep-seated cardiac, or arterial disease. The abdomen, like the chest, is in constant movement in connection with the act of respiration, and may be observed to be more prominent during inspiration, and more flat during expiration. These respiratory movements of the abdomen bear a certain relation to those of the chest, being often increased when the latter are arrested, and vice versa. Thus, in pleurisy, the respiratory movements are mostly abdominal, whereas in peritonitis, they are altogether thoracic. The variations observable in the disturbed relations of the respiratory movements in the thorax and abdomen are often highly instructive, especially in cases of dyspnæa from hydro-thorax, spasmodic asthma, anæmia, ascites, abdominal tumours, etc.

The inspection of deep-seated cavities, by means of specula, etc., belongs to the study of special diseases.

#### PALPATION.

Palpation also is a necessary mode of examination, and is sometimes practised by simply applying the tips of the fingers, at others by placing the hand on the part affected, and not unfrequently by employing both hands, and pressing with them alternately. This latter method is most applicable in endeavouring to judge of tumours, especially when large or deep seated, and situated in the abdomen. The position of the patient during palpation must be varied according to the part examined. The horizontal posture is

best to judge of deep-seated pulsations and vibrations, but sometimes the erect posture, or even leaning forward, becomes necessary, as when the heart is being examined. When feeling the abdominal organs through the integuments, these last should be relaxed, by causing some one to flex the inferior extremities on the abdomen, and push the head and neck forwards. In this manner palpation affords information—1st, As to the increased or diminished sensibility of various parts; 2d, Of their altered form, size, density, and elasticity; and, 3d, Of the different kinds of movement which they may be thrown into.


- 1. When pain is experienced in any part, it is generally increased by pressure and movement, if inflammatory, but is often relieved if neuralgic. Not unfrequently pressure causes pain or tenderness where otherwise neither are experienced. Thus deep pressure in the right iliac region causes pain in typhoid fever, which, however, must be judged of from the expression of the countenance, rather than the statement of the patient. Again, over the vertebral column pressure or percussion may induce pain that is otherwise not felt. In paralytic cases the diminution of sensibility can only be ascertained by feeling or pinching the part, and the limitation of anæsthesia is best arrived at by pricking the surface with some pointed hard substance.
- 2. Alteration in external form and size may be judged of by inspection, but in internal organs, especially abdominal ones, we derive more exact information from palpation conjoined with percussion, as in cases of hypertrophied liver and spleen, or when tumour exists. In such cases we can feel from the increased density and resistance the size and even outline of the morbid growths, which will be more or less distinct, in proportion as they are near the surface, and circumscribed in form. Occasionally organs are diminished in size, and cannot be felt in their normal positions, as when the inferior margin of the liver cannot be detected in this

way, from atrophy. The natural elasticity of parts may also be increased or diminished. Thus the abdomen is more elastic when air is in excess in the intestines, and less so when the peritoneum contains liquid. The integuments also may be more rigid and indurated, as in chronic skin diseases; or, on the other hand, soft and doughy, as in ædema, when they pit on pressure from diminished elasticity.

3. Certain motions in the thoracic and abdominal cavities, as well as in other parts of the body, are best judged of by palpation. It is in this way that the character and situation of pulsation at the heart, root of the neck, or elsewhere, are determined. The expansive motion of the thorax and abdomen during respiration is also thus ascertained. If we place the two hands, with the fingers spread out like a fan in the axillæ or flanks, and bring the two thumbs towards each other, near the sternum or umbilicus, we can judge by their approach and separation, the amount of expansion or retraction that takes place. Application of the hand also allows us to detect undulatory motions below the integument, and to determine the existence of vibrations, frictions, gratings, and crepitations. Rostan relates a case where all the symptoms of acute intercostal rheumatism were present (which disease was diagnosed), caused by a broken rib, that was overlooked from the diseased part not having been examined by palpation. There is a natural fremitus or thrill perceptible on placing the hand on the chest, when a person is speaking, which is increased in some diseases of the chest and lessened in others. This sensation is also sometimes felt in the large blood-vessels. It resembles more or less the vibration felt on placing the hand on the back of a cat while purring. Fluctuation is another sensation, caused by pressing on or percussing parts in such a way as to cause displacement of their contained fluids. A modification of it is known under the name of succussion, when produced by shaking the patient—a proceeding, however, which is seldom necessary.

#### MENSURATION.

The simplest way of measuring the circumference of parts, or the distance between any two fixed points, is by means of a graduated tape. In ascertaining the circular measurement of the chest or abdomen, that moment should



be chosen when the patient holds his breath at the end of an ordinary expiration, and great care should be taken that the tape is carried evenly round the body. The relative mensuration of the two sides of the chest or abdomen, is

Fig. 3. The chest-measurer of Dr. Sibson, natural size.

Fig. 4. The stethometer of Dr. Quain, one-third the real size.

best accomplished by choosing the spinous processes of the vertebræ as fixed points on the one hand, and a line drawn through the centre of the sternum and umbilicus on the other. The exact levels of the measurements should always be noted, such as at the nipples, margin of the lower ribs or




Fig. 5.

umbilicus, which are those most deserving of observation. The diameter of the trunk in various directions is best ascertained by means of a pair of callipers.

The amount of motion in the chest and abdomen, and of its various parts, are capable of being accurately determined by means of the chest measurer of Dr. Sibson (Fig. 3), or the stethometer of Dr. Quain (Fig. 4.) Both instruments are composed of a brass box, having a dial and index, which is moved by a rack attached to a prolonged pinion One revolution of the index indicates an or a string. inch of motion in the chest; the intervening space being graduated. It has been found necessary, when making observations on the respiratory movements, whether of the chest or of the abdomen, to divert the patient's atten-

Fig. 5. Mode of applying the stethometer when the string is used.—(After Quain.)

tion, and make him look straight forwards, otherwise these movements become so affected as to vitiate the results.



Fig. 6.

instruments may be applied in the sitting or recumbent pos-When a string is employed, as it may be with either instrument, the mode of application is represented

Fig. 6. Mode of applying the chest-measurer. - (After Sibson.)


Fig. 5. The method of using Dr. Sibson's chest-measurer by placing the pinion on the nail of the observer's left finger, moving with the chest, and steadying the right arm on the hip, or the hand on the back of a chair is seen, (Fig. 6).

The expansibility of the lungs, and the amount of air expelled from the chest after full inspiration, may also be measured by the spirometer of Mr. Hutchinson. But the necessity of determining the height and weight of the individual, of teaching him how to inspire and expire, of paying attention to the muscular force and other circumstances, so interferes with the correct conclusions to be derived from this mode of exploration, as to render it valueless in the examination of every case. But as a means of physiological research in determining the vital capacity of the chest, Mr. Hutchinson's investigations are of the utmost importance.

## LECTURE II.

#### PERCUSSION.

THE object of percussion is to ascertain the density and size of organs. It may be practised directly, or through the medium of an interposed body (mediate percussion). Without knowing how to strike, and to produce clear tones, we can never educate the ear, or the sense of resistance. This preliminary part in the art of percussion, requires a certain dexterity, which some find it very difficult to obtain. The difficulty seems to depend, in some cases, on a deficiency in the proportions usually existing between the length of



the fingers. Thus, I have seen more than one person who had the index finger nearly an inch shorter than the middle one, and who, consequently, found it impossible to strike the pleximeter fairly with the tips of two fingers at once. By far the most common cause of failure, however, is want of patience and perseverance in overcoming the first mechanical difficulties; and there is every reason to believe that

could this be surmounted, accurate percussion would become

more universal and better appreciated. Without entering into the numerous discus-

sions which have arisen as to the superior advantages of one plan as compared with another, or of using this or that instrument, I may mention, that for the last sixteen years I have employed a pleximeter and a hammer. These instruments I can confidently recommend to you as the readiest means of obtaining accurate results at the bed-side by means of percussion.

The ivory pleximeter I use is that of M. Piorry, as modified by M. Malliot. Its length is two inches, and breadth one. It possesses two handles, and an inch and a half scale drawn upon the surface. It may be applied with great precision to every part of the chest, even in emaciated subjects (Fig. 7).

The hammer is the invention of Dr. Winterich of Wurzburgh. The advantages it possesses are,—1st, That the tone produced, in its clearness, penetrativeness, and quality, far surpasses that which the most practised percussor is able to occasion by other means. 2d, It is especially useful in clinical instruction, as the most distant student is enabled to distinguish the varieties of tone with the greatest ease. 3d, It at once enables those to percuss, who,

Fig. 8. It at once enables those to percuss, who, from peculiar formation of the fingers, want of opportunity,

time, practice, etc., are deficient in the necessary dexterity (Fig. 8).

With the assistance of the instruments I now recommend to you, every student acquainted with the relative situations of the different thoracic and abdominal organs, is himself enabled, without other preliminary education, to detect the different degrees of sonoriety they possess in a state of health and disease. I may say, that by means of these instruments, after one hour's practice on a dead body, he is placed on a par (as regards the art of percussion) with the generality of experienced practitioners in this country; and any of you, after one month's employment of them, will be enabled to mark out accurately on the surface of the body, the size and form of the heart, liver, spleen, etc.

#### OF THE DIFFERENT SOUNDS PRODUCED BY PERCUSSION.

The sounds produced by percussion arise from the vibrations occasioned in the solid textures of the organ percussed. The different density and elasticity of these textures will of course more or less modify the number and continuance of the vibrations, and give rise to different sounds.

M. Piorry considers that nine elementary sounds are thus formed, which he has designated, from the organ or part which originates them, "femoral, jecoral, cardial, pulmonal, intestinal, stomachal, osteal, humorique, and hydatique." I consider that all these sounds may be reduced to three elementary ones; that, in point of fact, there are only three tones occasioned by percussion, and that all the others are intermediate. These three tones are respectively dependent,—1st, On the organ containing air; 2d, On its containing fluid; and, 3d, On its being formed of a dense uniform parenchymatous tissue throughout. These tones, therefore, may be termed the tympanitic, the humoral, and the parenchymatous. Percussion over the stomach gives

the best example of the first kind of sound; over the distended bladder, of the second; and over the liver, of the third. Certain modifications of these sounds occasion the metallic and the cracked-pot sound. The latter is made audible over the chest under a variety of circumstances, by percussing with the mouth open. The terms jecoral, cardial, pulmonal, intestinal, and stomachal, however, may be used to express those modifications of sound produced in percussing respectively the liver, heart, lungs, intestines, and stomach.

It must be remembered, however, that the tones even then may vary according to circumstances. Thus, immediately after a deep inspiration, the pulmonal sound will be rendered more tympanitic, and, after expiration, more parenchymatous. In the same manner, the stomach and intestines may give out different sounds according as they are more or less full of contents. In the left or right iliac fossa a clear tympanitic sound will be heard when the intestine below is empty, and a dull parenchymatous sound when it is full of foces.

A study of the different modifications of sound, which various organs thus produce in a state of health, readily leads to the comprehension of the sounds which may be elicited in a morbid state. Thus, the lungs may occasion a dull or parenchymatous sound, from solidification, the result of exudation, or, on the other hand, become more tympanitic, from the presence of emphysema. The abdomen may give out a parenchymatous sound, from enlargement of the uterus or an ovarian tumour; or a humoral sound, from the effusion of fluid in the cavity of the peritoneum.

OF THE SENSE OF RESISTANCE PRODUCED BY PERCUSSION.

By the sense of resistance is understood the peculiar sensation resulting from those impressions which are communicated to the fingers on striking hard, soft, or elastic bodies. It is of the greatest service in determining the physical condition of the organ percussed. The sense of resistance bears relation to the density of the object struck,—hence firm and solid textures offer more resistance than those which are soft or elastic. The thorax of the child is elastic, whilst that of the adult is unyielding. Of all the thoracic and abdominal organs, the liver presents the greatest degree of resistance, and the stomach the least. The presence of fluid in the hollow viscera, offers a medium of resistance between the parenchymatous organs on the one hand, and those containing air on the other. But air much condensed, or fluid contained within the rigid walls of the thorax, may offer a considerable degree of resistance.

The sense of resistance should be as much educated by the physician as the sense of hearing, and it would be difficult for an individual, practised in the art of percussion, to say which of these two points is the more valuable to him. Both are only to be learnt by practice, and considering it perfectly useless to describe that in words which may be learnt in half an hour, by the use of the pleximeter and hammer on a dead body, or the living subject, I shall now proceed to describe the

# GENERAL RULES TO BE FOLLOWED IN THE PRACTICE OF MEDIATE PERCUSSION.

1. The pleximeter should be held by the projecting handles between the thumb and index finger of the left hand, and pressed firmly down upon the organ to be percussed. Much depends upon this rule being followed, as the sound and sense of resistance are considerably modified according to the pressure made by the pleximeter. A very easy experiment will prove this. If, for instance, the pleximeter be struck while it rests lightly on the abdomen over the umbilicus, and again, when it is pressed firmly down

amongst the viscera, the change in tone will be at once perceived. In the first case, a dull sound is produced, from the muscles and integuments being alone influenced by the force of the blow; in the second case, a clear tympanitic sound is occasioned from the vibration of the walls of the intestine. In every instance, therefore, the pleximeter should be so held and pressed down, as to render it, so to speak, a part of the organ we wish to percuss.

- 2. Great care must be taken that no inequality exist between the inferior surface of the pleximeter and the skin. Firmly pressing it down will always obviate this when the abdomen is examined. As regards the thorax, the groove over the anterior mediastinum, the prominence of the clavicles and of the ribs, in emaciated subjects, may allow a hollow to exist under the instrument, by which a deceptive tympanitic sound is occasioned: By a little management, however, with the small and oval pleximeter I have recommended, this may readily be avoided.
- 3. The hammer should be held, as advised by Dr. Winterich, between the thumb and the first and third fingers, the extremities of which are to be placed in hollows prepared for them in the handle of the instrument. By some these are considered useless, but in all cases where slight differences in tone are to be appreciated, I have found this the best mode of employing it. Ordinarily, however, it will be sufficient to hold it by the extremity of the handle, merely in such a manner as will enable the practitioner to strike the pleximeter lightly, or with force, as occasion may require.
- 4. Care must be taken to strike the pleximeter fairly and perpendicularly. Unless this be done, vibrations are communicated to textures in the neighbourhood of the organ to be percussed, and fallacious results are the consequence. If in percussing the lungs, for example, the blow be made obliquely, we obtain the dull sound produced by the rib,

and I have seen considerable error in the diagnosis thus occasioned.

- 5. A strong or gentle stroke with the hammer will modify the tone and sense of resistance, inasmuch as the impulse may be communicated by one or the other to a deep-seated or a superficial organ. Thus a gentle stroke will elicit a pulmonal tympanitic sound just below the fourth rib, where a thin layer of lung covers the liver, but a strong one will cause a jecoral parenchymatous sound. At the inferior margin of the liver, on the other hand, where a thin layer of the organ covers the intestines, the reverse of this takes place, a gentle stroke occasioning a dull, and a strong one a clear sound.
- 6. By withdrawing the hammer immediately after the blow, we are better able to judge of the sound; by allowing it to remain a moment, we can judge better of the sense of resistance.
- 7. The integuments should not be stretched over the part percussed, as when the stethoscope is employed, for an unnatural degree of resistance is thus communicated to the hand of the operator from the muscular tension. In every case, especially where the abdomen is examined, the integuments and superficial muscles should be rendered as flaccid as possible.
- 8. It is always best to percuss on the naked skin. It is not absolutely essential, however; and in cases where, from motives of delicacy, it is desirable that the chest or abdomen be not exposed, it only becomes necessary that the covering of linen or flannel be of equal thickness throughout, and not thrown into folds.
- 9. The position in which the individual examined should be placed, will vary according to the organ explored. In percussing the thoracic organs and the liver, a sitting position is most convenient. The stomach, intestines, uterus, bladder, and abdominal tumours or effusions, are best exa-

mined when the patient is lying on the back, with the knees flexed so as to relax the abdominal walls, and, if necessary, the head and neck bent forward, and supported by pillows. In percussing the spleen, the individual should lie on the right side, and when the kidneys are examined he should lie on the breast and abdomen. In cases of effusion into the serous cavities, a change of position furnishes most valuable indications.

10. In percussing any particular organ, the pleximeter should be first applied over its centre, where the sound and sense of resistance it may furnish are most characteristic. Two blows with the hammer are generally sufficient to determine this. From the centre, the pleximeter should be moved gradually towards the periphery, or margin of the organ, and struck as it proceeds with the hammer, now forcibly, now lightly, until the characteristic sound of the next organ be elicited. The pleximeter is then gradually to be returned towards the organ under examination, until the difference of tone and sense of resistance become manifest. In this manner having first heard the two distinct sounds well characterised, we shall be better enabled to determine with accuracy the limit between the one and the other. This may be done exactly after having determined whereabouts the line of separation is, by placing the long diameter of the pleximeter transversely across it, and striking, first one end of the instrument, and then the other, till the precise spot is determined. This spot should now be marked, by placing with a pen a dot of ink on the skin. The opposite and then other portions of the margin of the organ should be limited in the same manner, and these in turn should be marked with dots of ink, until the whole organ be completely examined. Then by uniting all these dots with a line of ink, we have the exact form of the organ drawn upon the skin. When it is thought necessary to render the first mark permanent, in order to see if any subsequent change take place in the size of the organ, or extent of the dulness, it

may be rendered so, by carrying lightly a stick of argent. nit. over the ink line, while it is still moist.

Special Rules to be followed in percussing Particular Organs.

Before proceeding to percuss individual organs in persons labouring under disease, you should obtain a general




Fig. 9.

knowledge of the limits and intensity of dulness on percussing the thoracic and abdominal viscera in health. The accompanying figures convey this information with great accuracy, the depth of tint corresponding to the dulness of tone and amount of resistance. The normal sonoriety and dulness exhibited (Figs. 9 and 10) will enable you to compare with readiness the alterations revealed by percussion under a variety of diseased conditions.

No description will suffice to convey proper ideas of the various alterations of tone occasioned by percussing over the different thoracic and abdominal viscera. To become acquainted with these, it is absolutely necessary to apply the pleximeter to the body, and then half an hour's practice with this instrument and the hammer will be sufficient to render any one conversant with those which may be heard in a normal state.




Fig. 10.

## Lungs.—Percussion of the lungs generally bears refer-


Fig. 9, Anterior, and Fig. 10, posterior view of the limits and intensity of dulness on percussion. P, pulmonal sound; C, cardiac sound; H, hepatic sound; S, splenic sound; G, gastric sound (here the stomach is moderately distended with air); E, enteric sound. In the anterior view the intestines are tolerably free from air, except CO, colic sound, from distended colon. The descending colon and rectum are filled, and sound dull. HU, humoral sound, over a distended bladder; M, muscular, and O, osteal sounds.—(After Piorry.)

ence to a change in density, which is only to be detected by comparing the healthy with the morbid portions. The great practical rule here to be followed is, to apply the pleximeter to both sides of the chest in succession, with the same firmness, exactly in the same situation, and let the blow with the hammer be given with the same force. Care must be taken that the position of both arms be alike, as the contraction of the pectoral muscles on one side more than on the other may induce error. In short, every circumstance must be the same before it is possible to determine, in delicate cases, either from the tone or sense of resistance, whether change of density exist in the lungs. When circumscribed alterations are discovered in the pulmonary tissue, their limits may be marked out on the surface of the skin, in the manner previously indicated. In this way I have frequently succeeded in determining with accuracy the size and form of circumscribed indurations, arising from partial pneumonia and pulmonary apoplexy. Under the clavicles, the pleximeter must be applied with great firmness. Inferiorly, a thin layer of lung lies over the superior surface of the liver; and to determine the exact place where its inferior border terminates, the blows with the hammer should be very slight. Posteriorly, also, the pleximeter must be firmly applied, and the force of the blows considerable; but they should decrease in force inferiorly, where a thin layer of lung descends over the liver much deeper than anteriorly.

In a healthy state, a distinct difference may be observed in the sonoriety of the lungs immediately after a full expiration and a full inspiration. This does not take place when the tissue becomes indurated from any cause; and thus we are furnished with a valuable diagnostic sign. Congestion of the lung, and pneumonia in its first stage, cause only slight dulness and increased resistance, which, however, are readily detected by the practised percussor.

In the second and third stage of pneumonia, and in apoplexy of the lung, this dulness and resistance are well marked, and even an impression of hardness and solidity communicated to the hand. When, however, the lung is infiltrated with tubercle, the induration is most intense, and the greatest degree of resistance communicated.

Partial indurations from apoplexy or simple cancerous



and tubercular exudation, may be detected by percussion, even when deep-seated and covered by healthy portions of the lungs. In this case, by pressing with the pleximeter, and striking lightly, a tympanitic sound only is heard; but by pressing the pleximeter down firmly, and striking with

Fig. 11. Phthisis—Atrophied heart and liver—Prolonged abstinence. 1. Atrophied heart; 2, Infiltrated tubercle on left side; 3, The same on right side with a cavity; 4, Atrophied liver; 5, Spleen; 6, Unusual dulness over abdomen, from prolonged abstinence.—(After Piorry.)

force, the dull sound may be elicited and circumscribed. When indurations, however, exist inferiorly in those portions of the lungs which overlap the liver, it requires great practice to detect them with certainty. Caverns in the lungs, when large and filled with air, induce a tympanitic sound (Fig. 11, 3); but they are generally more or less full of viscous and fluid matters, and give rise to dulness.

Two or three ounces of fluid may be detected in the pleural cavity, by causing the patient to sit up. The height or level of the fluid is readily determined, and should be marked daily by a line made with nitrate of silver. If the




Fig. 12.

effusion be only on one side, the increased dulness is more easily detected. It disappears on placing the patient in

Fig. 12. Pleurisy. 1, On the right side when in the erect position; 2, On the left side, when lying on the right; 3, Kidneys; 4, Spleen.—(After Piorry.)

such a position as will cause the fluid to accumulate in another part of the pleural cavity, when the space which was previously dull becomes clear (Fig. 12). When the effusion entirely fills the pleural cavity, no limit, of course, can be detected; but, even then, the dulness is distinguished from that of the liver by the diminished feeling of resistance.

When the lung is emphysematous, or if air be present in the pleura, the sound becomes unusually tympanitic; this tympanitic note on percussion, however, may exist under a variety of circumstances, which it is of great importance to be acquainted with. Thus, condensation, from pneumonia at the posterior part of the lung, or partial pleurisy, by causing the anterior portion of the organ to be over-distended with air, or compressed and pushed forward, may give origin to this sound. The same occurs in chronic phthisis, over parts which were once dull, either from large dry cavities filled with air, or from the emphysema which accompanies cicatrices and partial condensation of pulmonary texture.

On percussing the chest with the mouth open, there may frequently be elicited a sound, which Laennec first likened to gently striking a cracked pot. It may be very closely imitated by crossing the palms of both hands, so as to leave a hollow between them, and then striking the knuckles of the inferior hand against the knee, so as to produce a clinking sound. I have produced it by percussing the chest in cases of pleurisy, pneumonia, and phthisis; of congested, apoplectic, and emphysematous lungs; and even when these organs were quite healthy, if, as in young subjects, the ribs were very elastic. The conditions which seem favourable for the production of this sound, are, 1st, A certain amount of confined or tense air in the tissue of the lung; 2d, Its sudden compression by a solid body; 3d, Communication of this air with the external atmosphere. Hence it is not

diagnostic of any particular disorder, or pathological state, such as a pulmonary cavity, so much as of a physical condition, which, however, if rightly interpreted, is likely to be of the utmost advantage in our efforts at detecting the nature of diseases.\*

Heart.—To mark out the precise limits of the heart, constitutes the first difficult lesson in the art of percussion. M. Piorry commences by determining the clear sound at the




Fig. 13.

upper end of the sternum, and bringing the pleximeter gradually downwards till the dull sound of the heart be heard.

\* See the author's "Clinical Investigation into the diagnostic value of the cracked-pot sound."-Edinburgh Medical Journal for March 1856.

Fig. 13. Pericarditis, pneumonia, and loaded rectum. 1, Pericarditis; 2, Pneumonia, separable from the extreme dulness of the liver; 3, Loaded rectum.— (After Piorry.)

I have found it best to place the instrument first under and a little inside the left nipple, where the cardiac dulness is most intense; then to carry it upwards, striking it continually with the hammer until the clear sound of the lung be elicited; then, by bringing it down again towards the heart, we shall readily distinguish the line where cardial dulness commences, and thus limit the superior margin of The same method is to be followed in determining the situation of the lateral margins, only carrying the pleximeter outwards or inwards, striking more and more forcibly with the hammer, until the clear tympanitic sound of the lung only be heard. It is more difficult to determine the situation of the apex of the heart; for as this rests on the diaphragm, and this again upon the left lobe of the liver, it cannot readily be distinguished from them. The size of the heart, however, may be pretty accurately estimated, by limiting its superior and lateral margins. In females, the left mammary gland should be drawn upwards and outwards by an assistant. In the natural position of the organ (Fig. 1) it is well to remember that the auricles are on the right, and the ventricles on the left side.

The normal size of the heart differs in different persons. As a general rule, however, it may be considered that, if the transverse diameter of the dulness measure more than two inches, it is abnormally enlarged. It has been known to measure seven inches. (Piorry.) In hydropericardium, the dulness has been remarked to exist rather at the superior part of the sternum, than on one side or the other. (Piorry, Reynaud.) In pericarditis it bulges out inferiorly (Fig. 13, 1). In hypertrophy and dilatation of the right auricle, the increased extent of the dulness stretches towards the median line, and sometimes passes over it (Fig. 14, 3). In similar hypertrophy of the left ventricle, the dulness extends on the left side more or less, according to the increased size of the heart (Fig. 14, 1, and Fig. 15). In concentric hypertrophy,

there is little or no enlargement, but the density is greatly increased, which is readily detected by the feeling of resistance.

The presence of tubercle in the lungs surrounding the heart; aneurisms or other tumours pressing upon or in the neighbourhood of the organ; hypertrophied liver, extensive empyema, etc. etc., may render its mensuration difficult or impossible. The changes in position of the heart from a pleurisy on one side pushing it towards the opposite one, or from the pregnant lineary, an ovarian tumour or ascites thrusting it upwards may also be determined by percussion, especially if the impulse can be distinguished by palpation or auscultation.

Liver.—Limitation of the size of the liver should be commenced by placing the pleximeter over the organ on the right side, where the dulness and resistance are greatest. It should then be carried upwards, until the clear sound of the lung be distinguished, when it ought again to be brought down, and the limit marked. This limit, however, may indicate either the inferior margin of the lung, or superior convex surface of the liver. Now, as a thin layer of lung descends in front of the liver, it will be necessary to determine where the tympanitic sound ceases inferiorly, by striking gently with the hammer, and where the parenchymatous sound ceases superiorly, by striking forcibly, so that vibrations may be communicated to the organ through the layer of lung. The space between these two lines thus marked on the surface is wider in some individuals than in others, and deeper and more extensive posteriorly than anteriorly. By carrying the pleximeter from the right side anteriorly, and then posteriorly towards the left of the patient, the whole superior margin may be thus detected, and marked with ink upon the surface, except where the liver comes in contact, through the medium of the diaphragm, with the apex of the heart. The inferior margin is for the most part readily detected. It must be remembered, however, that in the same manner as a thin layer of lung covers the upper margin, so a thin layer of liver descends on the right side over the intestine. It is, therefore, necessary to be cautious in deter-




Fig. 14.


mining the inferior margins, for a tolerably strong blow with the hammer may give rise to a tympanitic sound from the intestine, heard through the liver. The lower margin must be percussed in an inverse manner to the superior, and as we proceed downwards, the force of the blow should be diminished. The inferior margin of the liver is in general readily detected, from the contrast which, on percussion, its

Fig. 14. Hypertrophy of liver and heart. 1, Hypertrophied liver, which may be still further enlarged to the dotted lines over the abdomen; 2, Distended gall-bladder; 3, Hypertrophied right auricle—1, Hypertrophied ventricles; 4, Loaded cœcum; 5, Loaded rectum and descending colon.—(After Piorry.)

dulness and density present, contrasted with the tympanitic and elastic feel of the intestines and stomach.

The superior limit of this organ is generally found about two inches below the right nipple, or corresponding with the fifth rib. Its inferior border descends to the lower margin of the ribs. The extent of the jecoral dulness in the healthy state is in general two inches on the left side, three inches in the hepatic region anteriorly, and four inches in the hepatic region laterally. (Piorry.)

Variations in the size of the liver, from congestion,



inflammation, abscesses, hydatids, tumours, atrophy, etc. etc., may often be exactly determined by means of percussion. In icterus, the increase and diminution of this organ,

Fig. 15. Hypertrophied liver and spleen in leucocythemia—Enlarged heart.

1, Hypertrophied heart with dilatation; 2, Great dulness over the larger part of abdomen from enlarged liver on the right side; and enlarged spleen on the left.

—(Partly after Piorry.)


as evinced by lines marked on the skin, will generally be found to bear a proportion to the intensity of organic dis-When tumours are present, the inferior border often presents an irregular form. If the inferior lobes of the lung be indurated by tubercles or hepatisation, it becomes difficult or impossible to draw the limit between them and the liver. When fluid effusion exists in the pleura, the increased density of the liver still serves to distinguish it, through the humoral sound of the fluid; and, by changing the position of the patient, its upper edge in the majority of cases may be limited. In cases of ascites, we must lay the patient on the left side, in order to measure the right lobe - on the right side to measure the left lobe, and on the abdomen to percuss it posteriorly. Sometimes the right lobe of the liver is so enormously hypertrophied, that its inferior margin extends to the right iliac fossa (Fig. 15).

When the gall-bladder is much distended with bile, or contains gall-stones to any amount, it may readily be detected by percussion, and the dulness it occasions immediately under the inferior margin of the liver, anteriorly and somewhat laterally, be marked off (Fig. 14, 2).

Spleen.—In percussing the spleen, it is necessary that the patient lie on the right side, and it is advantageous that the examination be made before, rather than after, meals. Anteriorly the sonoriety of the stomach and intestines causes the margin readily to be distinguished. Posteriorly, however, where the organ approaches towards the kidneys, this is more difficult. Its superior and inferior margins may be made out by striking the instrument with some force, and following the rule (No. 10) previously given. This organ offers great resistance on percussion.

In health, the spleen never projects below the false ribs, even during a deep inspiration. Its general size is about four inches transversely and two inches deep. (Piorry.) In

diseased states it may be atrophied or enlarged. I have



seen it measure upwards of twelve inches long and eight wide, and it then may project upwards and downwards, as indicated by the dotted lines in Fig. 16. A pleuritic effusion, ascites, pneumonia, or tubercular deposition in the inferior lobe of the left lung, may render a limitation of this organ difficult or impossible. If the dulness cannot be detected, we may infer that its dimensions are small. (Mailliot.)

Stomach and Intestines.—The sounds elicited by percussion of the stomach and intestines are of the greatest service to the practitioner:
—1st, As furnishing him with the

means of determining the form of other organs, as the liver, spleen, or bladder; 2dly, As enabling him to distinguish the presence or absence of fœcal or alimentary matter; and, 3dly, As the means of diagnosing abdominal tumours. Hence it is incumbent on every physician to be able at once to recognise the difference between the tones furnished by the stomach, small and large intestines, under various circumstances. To arrive at this knowledge, it is necessary to be acquainted with the relative positions of the different abdominal viscera, and the regions of the abdomen to which they correspond. For instance, it is usually the liver and not the stomach that occupies the so-called epigastric region

Fig. 16. 1, Slightly enlarged spleen, pushed somewhat upwards. The dotted lines (2 to 6) indicate how the organ may be enlarged in various diseases. 7, Elongation downwards in leucocythemia.—(Slightly modified from Piorry.)

just below the end of the sternum. The last-named organ is for the most part situated within the left lower costal walls, just below the heart and the base of the left lung (Fig. 1.)

In exploring the abdomen by means of percussion, the pleximeter should first be placed immediately below the xiphoid cartilage, pressed firmly down, and carried along the median line towards the pubes, striking it all the way, now hard, now gently with the hammer. The different tones which the stomach, colon, and small intestines furnish will thus be distinctly heard. The pleximeter should then be carried laterally, alternately to the one side, and then to the other, till the whole surface be percussed. In this manner, the different tones produced by the cocum and ascending colon on the right side, as well as by the stomach and descending colon on the left, will be respectively distinguished from that furnished by the small intestines. sounds and sense of resistance will be modified according as the different viscera are full or empty, as any one can determine on his own body by means of the pleximeter and hammer. When the intestines are full of fluid or solid contents, such portions may be circumscribed and marked out on the surface of the skin. I have thus often succeeded in determining the internal margin of the colon, in its ascending, transverse, or descending portions. Sometimes a portion of intestine is found lying between the abdominal walls and the stomach. The latter, however, may be readily limited, by pressing down the pleximeter, causing the patient to eat or drink, or by examining after dinner. The small intestines are almost never deprived of the tympanitic sound—a circumstance by which they may readily be distinguished from the stomach and large intestines. The distance of any particular knuckle of intestine from the abdominal walls may be pretty accurately calculated by the force necessary to be employed in pressing down the pleximeter, and striking with the hammer, in order to elicit a tympanitic or dull sound.

It is unnecessary to point out the numerous circumstances, and morbid conditions, in which percussion of the abdomen may prove useful in practice. Displacements and variations in size of the stomach or intestines; femoral and scrotal hernia; mesenteric, ovarian, and other tumours; peritoneal adhesions and effusions-may all frequently be diagnosed, and their limits determined, by a careful examination with the pleximeter and hammer. By means of percussion, even the nature of the tumour may often be arrived at; as, for instance, whether it be fungus hæmatodes, scirrhous, encysted, osseous, etc., by the different degrees of resistance they possess. Care, however, must be taken not to confound with tumours an enlarged spleen or liver, a distended uterus or bladder, stomach full of alimentary matter, etc. It should also be remembered that when the patient lies on his back the percussion sound over the stomach is resonant, but when he stands it is generally dull from the gravitation of the food.

In a practical point of view, it is often useful to determine by means of percussion, whether an enema or a purgative by the mouth is likely to open the bowels most rapidly. If, for instance, there be dulness in the left iliac fossa, in the track of the descending colon, that part of the intestine must be full of feeces, and an enema is indicated. If, on the other hand, the left iliac fossa sound tympanitic, and the right sound dull, an enema is of little service, as it will not extend to the cocum, and purgatives by the mouth are indicated (Figs. 13 and 14).

The presence of fluid in the peritoneum may be determined with great exactitude by means of percussion, and the height of the fluid marked, as in the case of pleuritic effusion. In the same manner, a change of position furnishes similar results. Abdominal distension from accumulation of air

may also be determined. If it be within the intestine, the




Fig. 17.

tympanitic note is partial and limited, if in the peritoneal cavity more equable and diffused (Fig. 17).

Kidneys.—To percuss the kidneys, the patient should lie on the abdomen and chest; a position which allows any ascitic fluid that may be present to gravitate downwards, whilst the intestines float upwards. The dulness and great resistance offered by the renal organs are, under such circumstances, at once determined (Figs. 10 and 12). Their external margins may for the most be easily limited, in con-

Fig. 17. Dropsy of the abdomen, enlarged heart, and aneurism. 1, Aneurism projecting from the arch of the aorta on the right side; 2, Hypertrophied heart, especially of the right auricle; 3, Liver, pushed upwards; 4, Ascitic fluid, gravitating inferiorly, the patient being on the back; 5 and 6, Stomach and intestines, superiorly and anteriorly.—(After Piorry.)

sequence of the loud tympanitic note of the intestines, which can be elicited round their external circumference in the two flanks. Internally the dulness merges into that of the spinal column. Enlargement of one or both of these organs from calculous, or scrofulous nephritis, pyelitis, or other diseases, may in this manner be made out as seen (Fig. 12) on the left side. Atrophy of these organs is more difficult to determine with exactitude, but may be demonstrated by careful percussion.

Bladder.—This viscus is only to be detected by percussion, when it is more or less distended, and rises above the pubes. It may then be distinguished, and its circular margin limited, by observing the tympanitic sound of the intestines, on the one hand, and the dull humoral sound furnished by the bladder, with increased resistance, on the other. When covered by intestines, it will be necessary to press down the pleximeter with tolerable firmness, but not in such a manner as to give the patient pain. In the infant the situation of the bladder is not so deep in the pelvis, and a small quantity of fluid renders it cognizable by means of percussion.

A ready approximation of the state of the bladder will be found of great service in cases of fever, apoplexy, delirium, imbecility, paraplegia, etc. etc. In several cases it has been found dangerously distended, on percussing the abdomen to determine the state of the intestines.

I have here only noticed those circumstances in the art of percussion which may be readily accomplished, and which every one may master in a few months by care and attention. For a description of the more delicate points, such as percussion of the uterus and fœtus, accurately limiting the auricles and ventricles, determining and marking out the ascending and transverse portions of the arch

of the aorta, etc., I must refer you to the admirable works of MM. Piorry\* and Mailliot.†

\* De la Percussion Médiate, etc., Paris, 1828. Du Procédé Opératoire, Paris, 1831. De l'Examen Plessimètrique de l'Aorte Ascendante, et de la Crosse Aortique, etc. Archives Gén. de Méd. vol. ix., 1840, p. 431. On Percussion of the Uterus, and its Results in the Diagnosis of Pregnancy; Monthly Journal, 1846-7, p. 857. Atlas de Plessimètrisme, Paris, 1851.

† Mailliot (L.) Traité de la Percussion Médiate, etc., Paris, translated into English, with notes, by Dr. George Smith of Madras.

### LECTURE III.

#### AUSCULTATION.

THE object of auscultation is to ascertain and appreciate the nature of the various sounds which occur in the interior of the body. It has been found most useful when applied to the pulmonary and circulatory organs. Auscultation of the abdomen is occasionally serviceable, especially in certain cases of pregnancy, and during labour. It has also been applied to the head, although I have never been able to make out any useful results from the practice.

# GENERAL RULES TO BE FOLLOWED IN THE PRACTICE OF AUSCULTATION.

1. Auscultation may be practised directly by applying the ear to the part, or indirectly through the medium of a stethoscope. Generally speaking, direct auscultation answers every necessary purpose except when the surface is unequal, or when it is desirable to limit the sounds to a small region, as during auscultation of the heart. In either of these cases a stethoscope is necessary. The instrument is also useful to cenfirm or nullify the existence of certain fine sounds which may be detected by the naked ear; to remove the head of the practitioner a respectable distance from the bodies of persons not distinguished for cleanliness; and lastly, as the most delicate method of auscultating the chest anteriorly in women. You should regard the stetho-

scope as a mere means to an end—that end being, 1st, the education of the ear; and, 2d, a right appreciation of the pathological changes indicated by certain sounds.

- 2. In the choice of a stethoscope, you should observe, 1st, That the ear-piece fits your own ear; 2d, That the trumpet-shaped extremity is not above an inch and a half in diameter, and is rounded so as not to injure the patient's skin when pressure is made upon it; 3d, That it is light and portable. The instruments recently made of gutta percha fulfil all these conditions.
- 3. In applying the ear, the surface should be covered only with a smooth piece of linen or a towel. In using the stethoscope, it should be applied to the naked skin, and held steady immediately above the trumpet-shaped extremity by the thumb and index finger, pressed down with tolerable firmness, whilst the second, third, and fourth fingers enable you to ascertain whether the circular edge be perfectly applied, which is absolutely essential.
- 4. The position of the patient will vary according to the part examined. In auscultating the lungs anteriorly, the erect or recumbent positions may be chosen, when the two arms should be placed in a symmetrical position by the side. If the chest be examined posteriorly, the individual should lean somewhat forward and cross the arms in front. During auscultation of the abdomen, various positions will be required, according as the anterior, lateral, or posterior regions demand investigation. The practitioner, also, should choose such a position as will prevent too much stooping or straining. Generally speaking, the beds in the Infirmary here are too low, and render auscultation very fatiguing to the physician.
- 5. Whenever individuals are thrown into such a state of agitation as to interfere with the regular action of the heart or lungs, the examination should be deferred until their fear diminishes, or the greatest caution should be exercised in

drawing conclusions. Non-attention to this rule has led to many errors.

6. Before examining patients in a hospital, it is necessary that you should have made yourselves perfectly acquainted with the sounds which are continually going on in the healthy body. Omission of this rule not only renders the examination of patients useless, but betrays great want of consideration. For, as it is only from the alterations the healthy sounds undergo, or their being replaced by others, that we draw conclusions, how can this be accomplished if we are ignorant of their character in the first instance? It is expected, therefore, of every examining pupil, that he should be familiar with the character and theory of the various sounds heard in the healthy body before coming to the bedside. This study belongs to the Institutes of Medicine, rather than to that of Clinical Instruction.\*

## SPECIAL RULES TO BE FOLLOWED DURING AUSCULTATION OF THE PULMONARY ORGANS.

- 1. In listening to the sounds produced by the action of the lungs, we should pay attention to three things:—1st, The natural respiration; 2d, The forced or exaggerated respiration; and 3d, The vocal resonance. For this purpose, having listened to the sounds during ordinary breathing, we direct the patient to take a deep breath, and then, still
- \* Before commencing the actual examination of patients, you should make one or more serious, careful, and prolonged examinations of the chest of one of your fellow-students in private, so as to familiarise your ear with the healthy laryngeal, tracheal, pulmonary, and cardiac sounds, and with the character of the vocal resonance, and of the cough, as heard in various parts of the chest. You should then listen in the same manner to the chest of a young boy of from five to eight years of age, and observe how clear and exaggerated the pulmonary sounds are. Then read the short description of the healthy and morbid sounds which follow, and subsequently consult the more lengthened systematic treatises on this subject.

listening, we ask him a question, and during his reply judge of the vocal resonance.

- 2. You should commence the examination immediately under the centre of one clavicle; and having ascertained the nature of the sounds and vocal resonance there, you should immediately listen in exactly the corresponding spot on the opposite side. The examination should be continued alternately from one side to the other, in corresponding places, until the whole anterior surface of the chest is explored. The posterior surface is then to be examined in like manner.
- 3. When in the course of the examination, anything different from the normal condition is discovered at a particular place, that place and the parts adjacent should be made the subject of special examination, until all the facts regarding the lesion be ascertained.
- 4. It is occasionally useful to tell the patient to cough, in which case we are enabled to judge,—1st, Of forced inspiration, as it precedes the cough; and 2d, Of the resonance which the cough itself occasions.

# OF THE SOUNDS ELICITED BY THE PULMONARY ORGANS IN HEALTH AND IN DISEASE.

I am anxious to impress upon you, that the sounds which may be heard in the lungs are like nothing but themselves. Students are too apt to take up erroneous notions from reading on this subject, and, instead of listening to the sound actually produced, fatigue themselves in a vain endeavour to hear something like the crackling of salt, the rubbing of hair, foaming of beer, or other noises to which these sounds have been likened. Preconceived notions frequently oppose themselves to learning the truth, and have to be got rid of before the real state of matters can be ascertained. Hence the great importance of deriving your first impressions of the

sounds to be heard by auscultation, not from books or lectures, but from the living body itself.

If you listen through your stethoscope, placed over the larynx and trachea of a healthy man, you will hear two noises -one accompanying the act of inspiration, and the other that of expiration. These are called the laryngeal and tracheal sounds or murmurs. If you next place your stethoscope a little to the right or left of the manubrium of the sternum, you will hear the same sounds diminished in intensity. These are the bronchial sounds or murmurs. now you listen under and outside the nipple on the right side, or posteriorly over the inferior lobe of either lung, you will hear two very fine murmurs. That accompanying the inspiration is much more distinct than that accompanying the expiration. By some, on account of its excessive fineness, it is stated that there is no expiratory murmur in health; but this is incorrect. These sounds, then, are the vesicular respiratory murmurs. All these sounds become exaggerated during forced respiration, but in a state of health they never lose their soft character. Again, if you listen in the same places, whilst the individual speaks, you will hear a peculiar resonance of the voice, which has been called, in the first situation, pectoriloguy; in the second, bronchophony; while in the third it is scarcely audible. A knowledge of these circumstances, and a capability of appreciating these sounds, are necessary preliminary steps to the right comprehension and detection of the murmurs which may be heard during disease.

I have to suppose, then, that you have made your ears familiar with these sounds, and that you are acquainted with the present state of theory regarding their formation. This last may be stated in very few words to be, that the respiratory murmurs are occasioned by the vibration of the tubes through which the air rushes, according to well-known acoustic principles. Hence they are loudest in the trachea,

finer in the large bronchi, and finest in their ultimate ramifications. The vocal resonance, on the other hand, originates in the larynx; and diminishes or increases—1st, According to the distance of any point from the source of the sound; and, 2d, According to the power which the textures have in propagating it.

If now you examine, in succession, any six of the cases in the wards which are labouring under well-marked pulmonary diseases, you will have no difficulty in recognising that all the sounds you hear may be classified into two divisions:—1st, Alterations of the natural sounds; 2d, New, or abnormal sounds, never heard during health.

I. ALTERATIONS OF THE NATURAL SOUNDS.—All the sounds of which we have spoken, and which can be heard in the lungs during health, may, in certain diseased conditions, be increased, diminished, or absent; their character or position may be changed; and with regard to the respiratory murmurs, they may present alterations in rhythm or duration with respect to each other.

Alterations in Intensity.—Some persons have naturally louder respiratory murmurs than others; if this occur uniformly on both sides, it is a healthy condition. Occasionally, however, the sounds are evidently stronger in one place, or on one side (puerile respiration), generally indicating increased action of the lung, supplementary to diminished action in some other part. In the same manner, there may be feeble respiration simply from diminished action, as in feeble or old persons; but it may also be occasioned by pleurodynia, obstructions in the larynx, trachea, or bronchi—pleurisy, or pulmonary emphysema, or exudations filling up a greater or less number of the air-cells and smaller tubes, as in pneumonia, phthisis, etc. Complete absence of respiration only occurs where there is extensive pleuritic effusion hydrothorax or pneumothorax.

Alterations in Character.—The various respiratory murmurs may, in certain conditions of the lung, assume a peculiar harshness, which, to the ear of the practised auscultator, is a valuable sign, indicative of altered texture. Thus, in incipient phthisis, the vesicular murmur under the clavicle is often rude or harsh. In pneumonia the bronchial respiratory murmur presents a similar character. When ulceration exists, it becomes what is called cavernous (hoarse or blowing); and in certain cases of pneumothorax, it assumes an amphoric or metallic character.

Alterations in Position.—It frequently happens that the sounds which are natural to certain parts of the chest, are heard distinctly, where in health they are never detected. Thus, in pneumonia, bronchial, or tubular breathing, as it is sometimes called, may be evident, where only a vesicular murmur ought to exist. This is often well marked with regard to the vocal resonance, as certain lesions, which occasion condensation or ulceration, will enable us to hear either bronchophony or pectoriloquy in parts where, under ordinary circumstances, no voice can be heard.

Alterations in Rhythm.—In health, the inspiration is usually three times as long as the expiration. In certain diseased conditions this relation is altered, or even inverted. In incipient phthisis we often find the expiration unnaturally prolonged. In chronic bronchitis and emphysema it is three or four times longer than the inspiration.

- II. New or Abnormal Sounds.—These are of three kinds:—1st, Rubbing or friction noises; 2d, Moist rattles; 3d, Vibrating murmurs.
- 1. Rubbing or Friction Noises are caused in the pulmonary apparatus by some morbid change in the pleuræ, whereby, instead of sliding noiselessly on one another, they emit a rubbing sound. This may be so fine as to resemble the rustling of the softest silk, or so coarse as to sound like the

creaking of a saddle, grating, rasping, etc.; and between these two extremes you may have every intermediate shade of friction noise. This variation in sound is dependent on the nature of the alteration which the pleuræ have undergone. If covered with a softened thin exudation, the murmur will be soft; if it be tougher and thicker, the sound will be louder; if hard, dense, and rough, it will assume a creaking, harsh, or grating character, etc. etc. These noises are heard in the various forms of pleurisy.

- 2. Moist Rattles are produced by bubbles of air traversing or breaking in a somewhat viscous fluid. This may occur in the bronchi, when they contain liquid exudation, mucus, or pus, or in ulcers of various sizes. They may be so fine as to be scarcely audible (when they have been called crepitating), or so coarse as to resemble gurgling or splashing, when they have received the name of cavernous. Here again, between these two extremes, we may have every kind of gradation, to which auscultators have attached names, such as mucous, submucous, subcrepitating, etc. etc. With these names you need not trouble yourselves; all that it is important for you to determine is, that the sound be moist, and you will easily recognise that the rattles are coarse or large, in proportion to the size of the tubes or ulcers in which they are produced, and the amount of fluid present. These rattles may be heard in pneumonia, phthisis, pulmonalis, bronchitis, pulmonary apoplexy, etc. etc.
- 3. Dry Vibrating Murmurs arise when the air-tubes are obstructed, constricted, or lose their elasticity and become enlarged, whereby the vibrations into which they are thrown by the column of air, produce sounds or tones of an abnormal character. Hence murmurs may be occasioned of a fine squeaking (sibilous murmur), or of a hoarse snoring character (sonorous murmur), and between the two extremes, there may be all kinds of variations, to which ingenious people have applied names. These only cause confusion; all that

is necessary, being to ascertain that the murmur is dry, and you will readily understand that the fineness or coarseness of the sound will depend on the calibre of the tube or cavity thrown into vibrations. They are usually heard in cases of bronchitis and emphysema. Occasionally they present a blowing character, as when ulcers are dry, which often occurs in phthisis.

The vocal resonance, besides undergoing the changes already noticed in intensity, character, and position, may give rise to abnormal sounds. Occasionally it presents a soft reverberating or trembling noise, like the bleating of a goat (ægophony). The value of this sign, as indicative of pleurisy, was much overrated by Laennec. At present it is little esteemed. Sometimes the resonance gives rise to a metallic tinkling, a noise similar to that caused by dropping a shot into a large metallic basin, or the note produced by rubbing a wet finger round the edge of a tumbler or glass vessel. This is often best heard immediately after a cough in certain cas s of chronic phthisis. Egophony is supposed to be produced when a thin layer of serous fluid between the pleuræ is thrown into vibrations. The cause of metallic tinkling has created great discussion, and is not yet ascertained.

Such, then, are the principal sounds which may be heard by auscultation of the pulmonary organs in health and during disease. Many writers have endeavoured to point out their diagnostic importance, and drawn up rules which have always appeared to me much too arbitrary. Indeed, in so far as the education of medical students is concerned, I have long been persuaded that the study of these rules has retarded their powers of diagnosis, and afterwards led to dangerous errors in practice. I know of no dogma, for instance, more mischievous than the one which asserts a crepitating (that is a fine moist) rattle to be pathognomonic of

pneumonia, because it is just as common in phthisis, and is frequently heard in various other lesions of the pulmonary organs. Hence we should regard a crepitating rattle, not as distinctive of this or that so-called disease, but simply of fluid in the smaller air-passages;—and an increased resonance of the voice, as indicating hollow spaces with vibrating walls, or increased induration of the pulmonary textures, and not as diagnostic of phthisis, pneumonia, and so on. I wish, then, strongly to impress upon you,—

1st, That the different sounds are only indicative of certain physical conditions of the lung, and in themselves bear no fixed relation to the so-called diseases of systematic writers.

- 2d, No single acoustic sign, or combination of signs, is invariably pathognomonic of any certain pathological state,—and conversely, there is no pathological state which is invariably accompanied by any series of physical signs.
- 3d, Auscultation is only one of the means whereby we can arrive at a just diagnosis, and should never be depended on alone.

# SPECIAL RULES TO BE FOLLOWED DURING AUSCULTATION OF THE CIRCULATORY ORGANS.

- 1. In listening to the sounds produced by the action of the heart and arteries, we should pay attention—1st, To the impulse; 2d, To the character and rhythm of the sounds; 3d, To the place where they are heard loudest, and the direction in which they are propagated.
- 2. You should commence the examination by feeling for the spot where the apex of the heart beats against the walls of the chest, which will enable you to judge of the impulse. This ascertained, place your stethoscope immediately over it, and listen to the sounds. Then place the instrument above, and a little to the inside of, the nipple, near the

margin of the sternum, and listen to the sounds there. In the one situation you will hear the first or systolic sound, in the other the second or diastolic sound loudest.

- 3. If anything different from the normal condition be discovered in either one or the other position, or in both, they should be again carefully examined, and by moving the stethoscope below and round the apex of the heart, or above, in the course of the aortic arch or carotids, on the right and left side, etc. etc., it should be ascertained at what point, or over what space, the abnormal sounds are heard loudest, and whether they be or be not propagated in the course of the large vessels. Occasionally listening over the back and in the course of the descending aorta may be useful.
- 4. When, during the above examination, we discover a new source of impulse and of sound in one of the large vessels, this must be especially examined, the limits of such impulse and sound carefully ascertained—whether they be or be not synchronous with those originating in the heart—their direction, etc.
- 5. Under ordinary circumstances, the respiratory do not interfere with the detection of the cardiac sounds; but where the former are very loud and the latter indistinct, it is useful to direct the individual to hold his breath for a few moments. Sometimes the impulse and sounds of the heart are heard better by directing the patient to lean forward; they may also, if necessary, be exaggerated and rendered more distinct by directing him to walk up and down quickly, or to make some exertion for a short time.

OF THE SOUNDS ELICITED BY THE CIRCULATORY ORGANS IN HEALTH AND DISEASE.

On placing your car over the cardiac region in a healthy person, you will feel a beating, and hear two sounds, which have been likened to the tic-tac of a watch, but to which

they bear no resemblance. They may be imitated, however, very nearly, as pointed out by Dr. Williams, by pronouncing in succession the syllables, lupp, dupp. The first of these sounds, which is dull, deep, and more prolonged than the second, coincides with the shock of the apex of the heart against the thorax, and immediately precedes the radial pulse; it has its maximum intensity over the apex of the heart-below and somewhat to the inside of the nipple. The second sound, which is sharper, shorter, and more superficial, has its maximum intensity nearly on a level with the third rib, and a little above and to the right of the nipple-near the left edge of the sternum. These sounds, therefore, in addition to the terms first and second, have also been called inferior and superior, long and short, dull and sharp, systolic and diastolic-all which expressions, so far as giving a name is concerned, are synonymous.

The two sounds are repeated in couples, which, if we commence with the first one, follow each other with their intervening pauses, thus-1st, There is the long dull sound coinciding with the shock of the heart; 2d, There is a short pause; 3d, The short sharp sound; and 4th, A longer pause,—all which correspond with one pulsation. figures, the duration of these sounds and pauses by some have been represented thus,—the first sound occupies a third, the short pause a sixth, the second sound a sixth, and the long pause a third. Others have divided the whole period into four parts; of which the two first are occupied by the first sound, the third by the second sound, and the fourth by the pause. The duration, as well as the loudness of the sounds, however, are very variable even in health, and are influenced by the force and rapidity of the heart's action, individual peculiarity, and form of the thorax. Their extent also differs greatly. They are generally distinctly heard at the precordial region, and diminish in proportion as we withdraw the ear from it. They are less audible anteriorly on

the right side, and still less so posteriorly on the left side. On the right side posteriorly they cannot be heard. Their tone also varies in different persons; but in health they are free from a harsh or blowing character.

Great diversity of opinion has existed regarding the causes of these sounds-all of which you will of course have heard discussed before coming here. You must never forget, however, the cardiac actions which coincide with them; for our reasoning from any changes we may detect, will entirely depend upon our knowledge of these. We may consider, then, that there coincides with the first sound-1st, The impulse, or striking of the apex against the thoracic walls; 2d, Contraction of the ventricles; 3d, Rushing of the blood through the aortic orifices; and 4th, Flapping together of the auriculo-ventricular valves. There coincide with the second sound-1st, Rushing of the blood through the auriculo-ventricular valves; and 2d, Flapping together of the aortic valves. Contraction of the auricles immediately precedes that of the ventricles. The result of numerous pathological observations, and of many experiments, is, that in health the first sound is produced by the combined action of the auriculo-ventricular valves, of the ventricles, and of the rushing of the blood, which sound is augmented in intensity by the impulsion of the heart's apex against the thorax; whereas the second sound is caused only by the flapping together of the sigmoid valves.

With the cardiac as with the respiratory sounds, the alterations which take place during disease may be divided into—1st, Modification of the sounds heard in health; 2d, New or abnormal sounds.

I. Modifications of the Healthy Sounds.—These refer to the variations the healthy sounds present in their seat, intensity, extent, character, and rhythm.

Seat .- The sounds may be heard at their maximum in-

tensity lower than at the points previously indicated, as in cases of dilated hypertrophy of the left ventricle, enlargement of the auricles, or of tumours at the base, depressing the organ. They may be higher, owing to any kind of abdominal swelling pushing up the diaphragm. They may be more on one side or the other, in cases where the heart is pushed laterally by effusions of air or fluid in a pleural cavity. Various other circumstances may also modify their natural position, such as tumours in the anterior or posterior mediastinum, aneurisms of the large vessels, adhesions of the pericardium, deformity in the bones of the chest, etc. etc.

Intensity and Extent.—These are diminished in cases where the heart is atrophied or softened; when there is pericardial effusion, concentric hypertrophy of the left ventricle, or emphysema at the anterior border of the left lung. They are increased in cases of dilated hypertrophy, of nervous palpitations, and when neighbouring portions of the lung are indurated, especially in certain cases of pneumonia and phthisis pulmonalis.

Character.—The sounds become clearer or duller than usual, according as the walls of the heart are thinner or thicker. Occasionally they sound muffled in cases of hypertrophy or softening of the muscular walls. Not unfrequently there is a certain degree of roughness, which is difficult to determine as being healthy or morbid. Occasionally it ushers in more decided changes; at others, continues for years without alteration. These alterations in character are distinguished by some auscultators as variations in the tone of the sounds.

Rhythm or Time.—I need not say that the frequency of the pulsations differs greatly in numerous affections altogether independent of any special disease in the heart. In certain cardiac affections, however, the beats are intermittent, in others irregular—that is, they succeed each other at

unexpected intervals. The *number* of the sounds also varies. Sometimes only one can be distinguished, it being so prolonged as to mask the other. Occasionally three or even four sounds may be heard, depending either on reduplication in the action of the valves when diseased, or on want of synchronism between the two sides of the heart. Not unfrequently the increased and irregular movements of the organ, combined with the sounds, are of such a character as to receive the name of *tumultuous*.

II. NEW OR ABNORMAL SOUNDS.—These are of two kinds:—1st, Friction murmurs; 2d, Blowing or vibrating murmurs. Dr. Latham has called them exocardial and endocardial. I am in the habit of denominating them pericardial and valvular.

Pericardial or Friction Murmurs.—These murmurs are the same in character, and originate from the same causes, as the friction noises connected with the pulmonary organs. It is only necessary to observe, that occasionally they are so soft as closely to resemble blowing murmurs, from which they are only to be distinguished by their superficial character and limited extent.

Valvular or Vibrating Murmurs.—These murmurs vary greatly in character; some being so soft as to resemble the passage of the gentlest wind; others are like the blowing or puff from the nozzle of a bellows (bellows murmurs); whilst others are harsher, resembling the noise produced by grating, filing, sawing, etc. They are all occasioned, however, by diseases interfering with the functions of the valves. Sometimes these do not close, and the blood consequently regurgitates through them; at others, whilst this is the case, they are constricted, indurated, roughened, and even calcareous—whence the harsher sounds. They may be single or double, and have their origin either in the auriculo-ventricular or arterial valves, or in both at

once, the detection of which constitutes the diagnosis of the special diseases of the organ. Occasionally these sounds resemble musical notes, more or less resembling the cooing of a dove, singing or twittering of certain small birds, whistling, tinkling, etc. etc. These depend either upon excessive narrowing of the orifices, or upon any causes which induce vibrations of solids in the current of blood—as, when there are perforations in the valves, irregularities of their margins, string-like or other shaped exudations on their surface, etc. etc.

#### AUSCULTATION OF THE ABDOMEN.

On applying the stethoscope over the stomach and intestines in a healthy state, various gurgling and churning noises may be heard. In the former they may assume an amphoric or metallic character, in the latter they are called borborygmi. They are caused by the displacements of gas and water, and are most audible during the period of digestion, and the action of a purgative or enema. The impulse of the aorta can be detected, especially in thin subjects, when the pressure of the stethoscope may often be made to elicit a blowing sound.

In disease these sounds may be increased or diminished, and in addition, there may be present various kinds of friction or grating sounds when the surface of the peritoneum is roughened, owing to exudation or the unequal pressure of tumours. These last may also give rise to blowing murmurs, when it often becomes difficult to determine whether the morbid sound originates in the tumour itself, or from the pressure it exercises on the aorta. In cases of doubtful pregnancy, the marked rapidity of the fœtal pulse contrasted with that of the individual examined, constitutes a positive sign.

#### AUSCULTATION OF THE LARGE VESSELS.

On listening through the stethoscope placed over the arteries in the neighbourhood of the heart, we hear the same sounds as are produced at the sigmoid valves, propagated along its course, but more indistinct as we remove the instrument from the base of the heart. Those which are more distant have only one sound, which is synchronous with their impulse and their dilatation. This sound is of a dull character, but in health always soft.

In the various conditions of disease we have a single or double bellows sound, or it may be harsh, grating, rasping, etc. In the first place, you must ascertain whether any of these sounds are propagated along the artery from the heart, which you will know by listening over its course from that organ, and observing whether they increase as you proceed towards it. If the sound have an independent origin, it may originate from disease of the internal surface of the artery, when it will be harsh in proportion to the roughness; from stricture of, or pressure on the vessel, or from its dilatation. Generally speaking, the more dilated and superficially seated the vessel is, the sharper is the sound. Sometimes there is a double murmur in the course of a vessel, having an undoubted independent origin. is most common in cases where there is an aneurismal pouch, into which the blood passes in and out through an opening narrower than the swelling itself. Occasionally one or both such murmurs may possess somewhat of a metallic ringing, or even musical character, when the margins of the opening are probably tense, and thrown into peculiar vibrations.

Not unfrequently a soft systolic blowing is audible at the base of the heart, or over the carotids and deep jugular vein. At other times it is continuous, resembling humming, or the noise of a Parisian toy called *le diable*. These murmurs are distinguished from valvular ones—1st, By being systolic at the base of the heart; 2d, By their softness; 3d, By not being permanent; and, 4th, By occurring in anæmic or debilitated persons, and especially in young girls.

I have already told you never to form a conclusion from auscultation alone. Even when combined with percussion, it is not safe to form a diagnosis without a knowledge of all the circumstances of the case. Hence why I repudiate those rules which have been published in books, that have for their object the establishment of opinions from physical signs alone. At the same time, there can be no doubt that percussion and auscultation are absolutely essential to the proper investigation of maladies, although not more so than other modes of inquiry. I have, therefore, thought it best to give you a condensed resumé of the sounds which may be heard by auscultation of the lungs, heart, abdomen, and large vessels; pointing out a few of the diseased states in which they may be sometimes (not always) heard, and especially indicating the physical conditions on which they are supposed to depend. Their true diagnostic value can only be learned by the careful examination of individual cases.

I may conclude this subject by giving a few general rules diagnostic of pulmonary and cardiac diseases.

## GENERAL RULES DIAGNOSTIC OF PULMONARY DISEASES.

- 1. A friction murmur heard over the pulmonary organs indicates pleuritic exudation.
- 2. Moist, or dry râles, without dulness on percussion, or increased vocal resonance, indicate bronchitis, with or without fluid in the bronchi.
- 3. Dry râles accompanying prolonged expiration, with unusual resonance on percussion, indicate emphysema.
- 4. A moist râle at the base of the lung, with dulness on percussion and increased vocal resonance, indicates pneumonia.

- 5. Harshness of the inspiratory murmur, prolonged expiration, and increased vocal resonance confined to the apex of the lung, indicate incipient phthisis.
- 6. Moist râles, with dulness on percussion, and increased vocal resonance at the apex of the lung, indicate either advanced phthisis or pneumonia. The latter lesion commencing at or confined to the apex is rare, and hence these signs are diagnostic of phthisis.
- 7. Circumscribed bronchophony or pectoriloquy, with cavernous dry or moist râle, indicate a cavity. This may be dependent on tubercular ulceration, a gangrenous abscess, or a bronchial dilatation. The first is generally at the apex, and the last about the centre of the lung.
- 8. Total absence of respiration indicates a collection of fluid or of air in the pleural cavity. In the former case there is diffused dulness, and in the latter diffused resonance on percussion.
- 9. Marked permanent dulness, with increased vocal resonance and diminution or absence of respiration, may depend on chronic pleurisy, on thoracic aneurism, or on a cancerous tumour of the lung. The diagnosis between these lesions must be determined by a careful consideration of the concomitant signs and symptoms.

The general diagnostic indications now noticed as being derivable from physical signs, admit of several exceptions, which, however, it would be difficult to systematize. Hence they can only be acquired from a careful study of individual cases. It is important also to remember that these signs should never be relied on alone, but be invariably combined with a minute observation of all the concomitant symptoms. Thus the signs indicative of incipient phthisis may be induced by a chronic pleurisy confined to the apex, or by retrograde tubercle. In either case, the previous history, age, etc., may enable you to determine the nature of the lesion. Again, it may be impossible at the moment of examination

to distinguish between two diseases. For instance, there may be general fever, more or less embarrassment of the respiration, and pain in the side, accompanied with no dulness on percussion, but with a decided abnormal murmur, difficult to characterise, as being a fine moist rattle, or a gentle friction sound. Under such circumstances, the progress of the case will soon relieve you from any doubt as to whether a pleurisy or a pneumonia be present. The alterations which occur in the physical signs during the progress of the case also will indicate to the pathologist the changes which occur in the physical conditions and morbid lesions of the lungs. Thus the fugitive dry or mucous râles heard during a bronchitis, point out the occasional constrictions and obstructions in the bronchial tubes. The fine crepitation of incipient pneumonia, passing into absence of respiration, and this again into crepitation, will satisfy him as to effusion, solid coagulation, and subsequent softening of the exudation. In the same way, by an accurate appreciation of physical signs, and a thorough knowledge of morbid anatomy, the practised physician can tell the abnormal conditions produced by phthisis, pleurisy, etc., and judge from the symptoms the effects of these upon the constitution. with a degree of accuracy that to the tyro must appear to be marvellous. All such knowledge can only be acquired by constant examination of the patient on the one hand. and by a careful study of morbid anatomy in the pathological theatre on the other.

## GENERAL RULES DIAGNOSTIC OF CARDIAC DISEASES.

- 1. In health, the cardiac dulness on percussion measures two inches across, and the extent of dulness beyond this measurement indicates either the increased size of the organ, or the extent of pericardial effusion.
- 2. In health, the apex of the heart may be felt and seen to strike the chest between the fifth and sixth ribs, imme-

diately below and a little to the inside of the left nipple. Any variations that may exist indicate the altered positions of the apex in disease.

- 3. A friction murmur indicates pericardial exudation.
- 4. A bellows murmur with the first sound, heard loudest over the apex, indicates mitral insufficiency.
- 5. A bellows murmur with the second sound, heard loudest at the base, indicates aortic insufficiency.
- 6. A murmur with the second sound, loudest at the apex, is very rare, but when present it indicates—1st, Aortic disease, the murmur being propagated downwards to the apex; 2d, Roughened auricular surface of the mitral valves; 3d, Mitral obstruction, when it is almost always associated with insufficiency, and the murmur is double.
- 7. A murmur with the first sound, loudest at the base, and propagated in the direction of the large arteries, is more common. It may depend,—1st, on an altered condition of the blood, as in anæmia; 2d, on dilatation or disease of the aorta itself; and 3d, on stricture of the aortic orifice,—in which case it is almost always associated with insufficiency, and the murmur is double.
- 8. Hypertrophy of the heart may be independent of valvular disease, but this is very rare. In the vast majority of cases it is the left ventricle which is affected, in connection with mitral or aortic disease. In the former case the hypertrophy is equal with rounding of the apex; in the latter there is dilated hypertrophy with elongation of the apex.

Attention to these rules alone will, in the great majority of cases, enable you to arrive with precision at the nature of the lesion present. In cases in which there may be any doubt, you will derive further assistance from an observation of the concomitant symptoms, such as,—1st, the nature of the pulse at the wrist; 2d, the nature of the pulmonary or cerebral derangements. Thus, as a general rule, but one on which you must not place too much confidence, the pulse

is soft or irregular in mitral disease, but hard, jerking, and regular in aortic disease. Again, it has been observed that cerebral symptoms are more common and urgent in aortic disease, and pulmonary symptoms more common and urgent in mitral disease.

I purposely say nothing of diseases of the right side of the heart, and of a few other rare disordered conditions of the organ,—1st, because they occur so seldom as scarcely to merit our attention; and, 2d, because I am convinced that an appreciation of the rules above given is the best method of enabling you to comprehend and easily detect any exceptional cases which may arise. In truth, however, we have seen in our examinations at the bed-side, that your difficulty is, not how to arrive at correct conclusions from such and such data, but how to arrive at the data themselves. You have to determine,-1st, by percussion, whether the heart be of its normal size or not; 2d, whether an abnormal murmur does or does not exist; 3d, if it be present, does it accompany the first or second sound of the heart; and 4th, at what place and in what direction the murmur is heard loudest. These points ascertained, the conclusion flows from the rules previously given. But no instruction on my part, no reading or reflection on yours, will enable you to ascertain these facts for yourself. In short, nothing but percussing the cardiac region with your own hands, and carefully listening to the sound with your own ears, can be of the slightest service, and the sooner you feel convinced of this truth the sooner are you likely to overcome these preliminary difficulties.

## LECTURE IV.

### THE USE OF THE MICROSCOPE.

A KNOWLEDGE of the ultimate structure of the human body in its healthy and diseased conditions, is now so advanced as to necessitate the introduction of the microscope among the instruments of the medical practitioner. You must not suppose that an additional method of gaining information implies abandonment of those, the utility of which has stood the test of experience. Men must learn the everyday use of their senses; must know how to feel, hear, and see, in the same manner as they did before instruments were invented. We don't see the stars less clearly with our naked sight, because the telescope is necessary for an astronomer. Neither should a physician observe the symptoms of a disease less accurately because he examines the chest with a stethoscope, or a surgeon be less dexterous with the knife, because it is only by means of the microscope he can determine with exactitude the nature of a tumour. But it is unnecessary to enter into a lengthened argument to prove that the science and art of medicine are greatly indebted, in modern times, to the invention and proper application of ingenious instruments. The following examples will serve to convince you that the microscope is one of these :--

Example 1.—Some years ago I was summoned to see a

dispensary patient labouring under bronchitis, who was spitting florid blood. On examining the sputum with a microscope, I found that the coloured blood corpuscles were those of a bird. On my telling her she had mixed a bird's blood with the expectoration, her astonishment was unbounded, and she confessed that she had done so for the purpose of imposition.

Example 2.—A gentleman, for some years, had laboured under a variety of anomalous symptoms, referable to the head and digestive systems, under which he had become greatly reduced. He had consulted many practitioners, and visited innumerable watering places, in a vain search after health. On examining the urine with a microscope, I found it crowded with spermatozoa. He evidently laboured under spermatorrhæa, a disease which had never been suspected, but which was readily cured on the employment of an appropriate treatment.

Example 3.—A boy was brought to me with an eruption on the scalp, which was of so indefinite a character that its nature could not be determined. He had lately been elected to occupy a vacancy in one of our charitable educational establishments, and the question to decide was, whether the disease was or was not contagious. On examining the scab with a microscope, I readily discovered the Achorion Schoenleini, or fungus constituting true favus; and as this has been experimentally proved to be inoculable, I had no hesitation in preventing his admission to the school.

Example 4.—A child was supposed to be affected with worms, because it passed in abundance yellowish shreds, which, to the naked eye, closely resembled ascarides. All kinds of vermifuge remedies had been tried in vain. On examining the shreds with a microscope, I found them to

consist of the undigested spiral vessels of plants; and they ceased to appear when the vegetable broth used as food was abandoned.

Example 5.—I was called to see an infant, a month old, which was in a state of considerable emaciation, with constant diarrhea. The mother, however, maintained that her milk was abundant, and that it was taken in sufficient quantity. On being examined with a microscope, it was found to contain numerous granular cells and masses with comparatively few milk globules. In short, it presented, in an exaggerated degree, all the characters of colostrum, and this thirty days after delivery. It was evident, then, that the quality of the milk was in fault, an opinion which was confirmed by the recovery of the infant, when a healthy nurse was procured.

Example 6.—An individual was supposed to be labouring under dysentery, from the frequent passage of yellowish pulpy masses in the stools, accompanied with tormina and other symptoms. On examining these masses with the microscope, I found them to consist of undigested potato skins. On inquiry, it was ascertained that this person had eaten the skins with the potatoes. On causing these to be removed before dinner, the alarming appearance ceased, and the other symptoms also disappeared.

Example 7.—An elderly lady conceived herself to be affected with insects continually forming in the skin, which produced incessant itching and tingling. All the hair was removed, and every kind of application, including mercurial preparations, were tried without effect. On rubbing the surface, she always saw minute white rolls and black specks, which she regarded as insects in different stages of development. The torment and anxiety this caused her

for many months it is scarcely possible to conceive. At length she laboured under the idea that she was communicating the disease to her husband and daughter, when, at the request of her medical attendant in the west of Scotland, she came to Edinburgh, in order that I might investigate and treat it. I had the pleasure of showing this lady, under the microscope, that the white bodies were minute rolls of epidermis or of the cotton cloth with which she rubbed the skin, and that the black specks were portions of dust or soot. Her hallucination being in this way dissipated, she returned home perfectly well.

Example 8.—A child had been suffering for four years from copious and fœtid discharge from the nostrils, accompanied with great pain. At the end of that time, a dark brown and indurated mass was discharged about an inch long, and a quarter of an inch broad, closely resembling a sequestrum of bone. This mass I was requested to examine microscopically by Dr. Littlejohn, whose case it was, and from its structure I readily determined that it consisted of some fir wood. When this was known, the parents remembered that, about the time the disease commenced, alterations were made in the house, and that the children used to play with the wood shavings. There could be little doubt that a piece of shaving had been thrust up the nose, and been the cause of all the symptoms.

Examples of this kind could be readily multiplied. No doubt mistakes will be made with this instrument in the hands of inexperienced persons, in the same manner as the use of the stethoscope, or of a scalpel, may lead to a false conclusion, or to an accident. But this, so far from being an argument opposed to their employment, only proves the necessity of becoming more skilful in their use. Certainly there is none which requires more expert management in

itself, or more caution in drawing conclusions from its employment, than the microscope.

### DESCRIPTION OF THE MICROSCOPE.

It is not my intention to enter upon a description of the optical principles on which microscopes are constructed,




Fig. 18.

although you will find a knowledge of these very useful. I shall suppose that you are desirous of obtaining an instrument that will answer all the purposes of the anatomist and physiologist, as well as afford you every possible assistance in the way of diagnosis as medical men. For this purpose, you should learn to distinguish what is necessary from what is unnecessary, in order that you may procure the former in as convenient a form, and at as moderate a cost as possible.

A microscope may be divided into mechanical and optical parts. The former determine its general form and appearance. Of the numerous models which have been invented, the one here figured, exactly one-fourth its real

Fig. 18. Oberhaeuser's model, made at my suggestion, for medical men. One-fourth the real size.

size, appears to me the most useful for all the purposes of the physiologist and medical practitioner. The body consists of a telescope tube, eight inches in length, held by a split tube, three inches long. It may be elevated and depressed with great readiness by a cork-screw movement communicated to it by the hand, and this constitutes the coarse adjustment. It is attached to a cross bar and pillar, at the lower portion of which last, very conveniently placed for the hand of the observer, is the fine adjustment. The stage is three inches broad, and two and a half inches deep, strong and solid, with a circular diaphragm below it. The base of the instrument is heavily loaded with lead to give it the necessary steadiness.

This form of microscope possesses all the mechanical qualities required in such an instrument. These are—1st, Steadiness; 2d, Power of easy adjustment; 3d, Facility for observation and demonstration; and, 4th, Portability.

- 1. Steadiness.—It must be evident that if the stage of the microscope possesses any sensible vibration, minute objects, when magnified highly, so far from being stationary, may be thrown altogether out of the field of view. Nothing contributes more to the comfort of an observer than this quality of a microscope, and great pains have been taken to produce it. In the large London instruments this end has been admirably attained, but at so much cost and increase of bulk as to render it almost useless. In the small model I have recommended, all the steadiness required is present in the most convenient form.
- 2. Power of Easy Adjustment.—It is a matter of great importance to those who use the instrument much, and work with it for hours together, that the adjustments should work easily and rapidly, and be placed in convenient situations. Nothing can be more commodious than the man-

ner in which these ends are arrived at in the model figured. By insertion of the body of the instrument within a split tube, you may by a spiral movement, elevate and depress it with the greatest rapidity, and even remove it altogether if necessary. The necessity of continually turning the large screws affixed to most microscopes, becomes fatiguing in the extreme. Then the fine adjustment, placed conveniently behind the microscope, near the hand which rests on the table, is in the very best position; whereas, in some London instruments, it is placed on the top of the pillar, so that you must raise your hand and arm every time it is touched. In other London instruments, it is placed in front of the body, so that you must stretch out the arm and twist the wrist to get at it. No one could work long with so inconvenient a contrivance.

3. Facility for Observation and Demonstration.—For facility of observation and demonstration, it is necessary that the instrument should be of a convenient height, and that the stage on which the objects are placed should be easily accessible. Here, again, nothing can be more commodious than the microscope I have recommended, for, when it is placed on a table, its height is almost on a level with the eye, and we can look through it for hours without the slightest fatigue. On the other hand, the stage is elevated, just so much as enables the two hands, resting on their external edges, to manipulate with facility all kinds of objects placed upon it. The large London instruments are so high as to render it necessary to stand up to see through them. To obviate this disadvantage, a movement is given to the body, by which it can be depressed to any angle. But this movement renders the stage-oblique, and removes it to a distance, where it becomes very inconvenient to manipulate on its surface. To obviate this difficulty, the stage itself has been rendered movable in various ways by different screws, so that in this way complexity has been added to complexity, until a mass of brass work and screws is accumulated, to the advantage of the optician, but to the perplexity and fatigue of the observer. But by no contrivance is it possible to avoid the aching arms which such a position of the stage invariably produces in those who work with such a cumbrous machine for any length of time.

4. Portability.—This is a property which should by no means be overlooked in instruments that are intended more for utility than ornament. A medical man is often called upon to verify facts in various places; at his own house, at an hospital, at the bed-side of his patient, or at a private post-mortem examination. It is under such circumstances that the value of portability is recognised. The large London instruments require an equipage or a porter to transport them from place to place; even the putting them in and out of the large boxes or cabinets that are built around them, is a matter of labour. In short, notwithstanding the splendour of the screws, the glittering of the brass, and the fine workmanship, there can be little doubt that, on the whole, they are very clumsy affairs.

There are many occasions on which a medical man may find it useful to carry a microscope with him, especially in the case of post-mortem examinations. Many attempts have been made to construct a pocket microscope; and for the purposes above alluded to, I myself caused one to be constructed some years ago, which, with its case, resembled a small pocket telescope. Dr. Gruby of Paris, however, has planned the most ingenious instrument of this kind, which possesses most of the properties we have enumerated, and will be found very useful for those accustomed to microscopic manipulation. It is contained in a case, the size of an ordinary snuff-box, and possesses all the conveniences of the larger instruments, with various lenses, a micrometer,

slips of glass, needle, knife, and forceps, in that small compass. The prefixed woodcuts, exactly one-half the real




Fig. 19.

size, will give an idea of this ingenious microscope, manufactured by Brunner of Paris. For a more minute description, I must refer you to the "Monthly Journal of Medical Science" for December 1846.

There is a general feeling among the public that the larger a microscope is, the more it must magnify; but I need not tell you this is error. A very imposing mass of brass work and mechanical complexity, is no guarantee that you will see objects better, or, what is of more consequence, become good observers. On the contrary, the more unwieldy the instrument, the less disposed will you be to use it. Besides, the habitual employment of artificial methods of moving about the object, as by the screws of a moveable stage, will prevent your acquiring that dexterous use of your fingers and accuracy of manipulation which are at all times so useful. Nothing indeed, can be more amusing than

Fig. 19. Gruby's compound pocket microscope, exactly one-half the real size.

to see a man twisting his screws, pushing his heavy awkward stage about, and laboriously wasting time to find a minute object which another can do in a moment, and without fatigue, by the simple use of his fingers. But perhaps you





Fig. 20.

will consider the weightiest objection to the large instruments is the expense they necessitate,—the cost being necessarily in proportion to the amount of brass and mechanical labour employed upon them. If, then, you have to choose between a complex model and a simple one, I strongly advise you, as a matter of real economy, to choose the latter. Indeed the former, to a practical histologist, is worthless.

A very simple form of miscroscope was presented to the Physiological Society of Edinburgh, at its meeting on the 7th of June 1851, by Dr. W. T. Gairdner (Figs. 21 and 22). It consists of a Coddington lens, so small that the focal distance

Fig. 20. The same microscope mounted, ready for use.

is about the  $\frac{1}{15}$ th of an inch, and the magnifying power from 150 to 250 diameters linear. This lens is fixed in a round



plano-concave disc of brass, attached to a small handle of brass. On the plane side is a ring of silver, in which a thin piece of glass is fitted, also supported by a handle of brass. The two handles are united together by means of a fine screw, so that exact focal distance is attained. A drop of fluid placed on the *outside* of this glass, either covered or not with another glass, then applied to the eye and directed towards the light, will enable us to distinguish blood, pus, epithelial, or other corpuscles, various forms of crystals, etc.

GAIRDNER'S SIMPLE CLINICAL MICROSCOPE.—Fig. 21, a lateral view of the instrument. Fig. 22, a front view, showing in outline the posterior glass separated and turned aside.

sufficient, to experienced eyes, for the purposes of diagnosis. By shading the lens externally with the finger, all the effects of a diaphragm can be produced. It must not be supposed that this instrument will ever supersede the necessity of studying histology by means of a larger one; but, to him who is already familiar with minute objects, it will prove a valuable acquisition at the bedside. It is made by Bryson, optician, Princes Street.

We have next to speak of the optical parts of microscopes, which are certainly much more important than the mechanical ones,—everything depending upon obtaining a clear and distinct image of the object examined. Under this head we may describe the objective, the eye-piece, and methods of illumination.

1. The Objective, or series of Achromatic Lenses, is that part of the optical portion of a microscope which is placed at the bottom of the tube or body, and is near the object to be examined. This may be considered the most important part of the instrument, and the greatest pains have been taken by all opticians in the manufacture of good lenses. It is here I consider that the London opticians are pre-eminent, for I am not aware that in any part of the world such perfect objectives have been manufactured as the eighth of an inch by Smith, the twelfth of an inch by Ross, and the sixteenth of an inch by Powell. But when we come down to the one-fourth of an inch, which is by far the most useful objective for an atomical and medical purposes, the superiority of the London opticians is very slight, if any. At this magnifying power the compound lenses of C. Chevalier, Oberhæuser, Brunner, and Nachet of Paris, Schiek and Pistor of Berlin, Frauenhofer of Munich, and Ploesl of Vienna, may be employed with the greatest confidence, and it may be said, that by far the largest number of important discoveries in science have been made through their employment. The

Parisian lenses, in addition, have one great advantage, namely, their cheapness.

The London opticians have succeeded in combining the lenses of their objectives, so as to obtain a large field of vision, with as little loss of light as possible. These qualities are valuable in the lower magnifying lenses during the examination of opaque objects, and in the higher ones when observing transparent objects by transmitted light. But in the lenses of medium power, such as the one-fourth of an inch, the amount of light is so great as to be almost a defect. Notwithstanding careful management of the mirror and diaphragm, the field of vision is often dazzling, and always presents a glare most detrimental to the eyes of the observer. I cannot employ their fourth of an inch for fifteen minutes without feeling intense headache, and I know of more than one excellent observer in whom the sight has so much suffered from this cause as to incapacitate them from continuing their researches. In the same manner, the lenses of Brunner and Nachet give rise to a yellow light highly disagreeable; while those of Oberhaeuser, Schiek and Pistor, and Frauenhofer (with Amici's and Ploesl's I am not familiar), present a pale blue light, most pleasant to work with, and which may be gazed at for hours without fatiguing the eye.

For the above reasons, as well as from considerable experience in the use of many kinds of microscopes by different manufacturers, I am satisfied that the best lens you can employ for ordinary purposes is Oberhaeuser's No. 7, which corresponds to what is called in England the quarter of an inch. For low powers you may have Oberhaeuser's No. 3, or the one inch lens of the London opticians. For all the wants of the medical man these will be sufficient. The anatomist may occasionally require a higher lens, as during the examination of the ultimate fibrillæ of muscle, when the eighth, twelfth, or sixteenth of an inch of the London opticians

may be procured. All these lenses may be attached to the model we have recommended by means of a brass screw made on purpose.

- 2. The Eye-piece.—This is that portion of the optical apparatus which is placed at the upper end of the tube or body, and is near the eye of the observer. While the objective magnifies the object itself, the eye-piece only magnifies the image transmitted from below. Hence, as a source of magnifying power, it is inferior to the lens; and when this possesses any defects, these are enlarged by the eye-piece. Two eye-pieces are all that is necessary with the model I have recommended, and those of Oberhaeuser, called Nos. 3 and 4, are the most useful for the medical man.
- 3. Methods of Illumination.—There are few things of more importance to the practical histologist than the mode of illumination. This is accomplished—1st, By transmitted light; 2d, By reflected light; and 3d, By achromatic light.

Transmitted light is obtained by means of a mirror, placed below the object, which, to be seen, must therefore be transparent. In large microscopes the mirrors are provided with universal joints, so that they may easily be turned in any direction. Below the stage every microscope should possess a diaphragm pierced with variously sized holes, whereby the amount of light furnished by the mirror may be moderated. In Oberhaeuser and Nachet's instruments the smallest aperture should be employed for the higher objective. It is also useful in the examination of many objects that the light should be directed upon them sideways; this may be done by the diaphragm or by the mirror, and, in the small model formerly figured, is admirably attained by simply turning the whole microscope. The best light for microscopic purposes is that obtained by catching the rays which are reflected from a white cloud.

The conjoined use of the mirror and diaphragm can only be learned from actual experience.

Reflected light is employed in the examination of opaque objects, and the lenses of low power, manufactured by the principal London opticians, enable us to do so without assistance. Occasionally, however, the light of the sun is useful; and when this cannot be obtained, the rays of a lamp or gas light, concentrated by a bull's-eye lens, may be employed. Hence every microscope should be possessed of such a lens, and it is most convenient to have it attached to the body of the instrument by a moveable ring, and stem with two joints, as in the model figured, p. 86.

Achromatic light is only servicable in the examination of very delicate objects, with high powers. The apparatus necessary for obtaining it is occasionally useful in ascertaining the ultimate structure of muscle, or the nature of the markings on minute scales or fossils, but is useless for the purposes of the medical man. In the same way I know of no benefit to be obtained by a polarising apparatus.

In addition to the mechanical and optical parts constituting the microscope itself, the box which contains it should possess a convenient place for holding a few slips of glass, a pair of small forceps, a knife, and two needles firmly set in handles. A micrometer to measure objects with is also essential to those who are making observations with a view to their exact description. No other accessories are necessary.

An excellent microscope of the model previously figured, by Oberhaeuser (Fig. 18), with two objectives (Nos. 3 and 7), two eye-pieces (Nos. 3 and 4), a neat box with all the accessories necessary (with the exception of a micrometer, which had better be English) may be obtained in Paris for the sum of about 150 francs (£6), and will cost in Edinburgh, after payment of carriage and duty, about seven guineas. Nachet

and Brunner's instruments are much cheaper, as are the smaller models of Oberhaeuser. Either of them, for all the purposes of the medical man, is amply sufficient.

Test-Objects.—The defining power of a microscope is generally tested by examining with it a transparent object, having certain fine markings, which can only be rendered clearly visible when the glasses are good. In all such cases, it is of course necessary to be familiar with the structure of the test-object in the first instance. If you are not confident on this point, it is better to trust to the judgment of a friend, whose knowledge of histology is ascertained, or place your dependence entirely on a respectable optician. One of the best test objects for a quarter of an inch lens is a drop of saliva from the mouth. For, if the microscope shows with clearness the epithelial scales, the structure of the salivary globules, their nuclei, and contained molecules, you may be satisfied that the instrument will exhibit all the facts with which, as medical men, you have to do.— (See Fig. 24.)


## MENSURATION AND DEMONSTRATION.

Having, then, obtained a good instrument, and tested its qualities in the manner described, you should next determine the number of diameters linear the various combinations of glasses magnify. This you may do for yourself with the aid of a micrometer, a pair of compasses, and a measure.

A micrometer is a piece of glass on which lines are ruled at the distance of 1-100th or 1-1000th of an inch. This must be placed under the instrument, when the lines and the distances between them will of course be magnified by the combination of glasses employed, like any other object. Taking a pair of compasses in one hand, we separate the points, and place them on the stage (always on a level with

the micrometer magnified). Now, looking through the instrument with one eye, we regard the points of the compasses with the other, and mark off by the naked sight, say the 1-100th of an inch, as magnified by the instrument. Though difficult at first, a little practice enables us to do this with the greatest accuracy. The result is, that if the distance magnified and so marked off (1-100th of an inch) is equal to three inches, the instrument magnifies 300 times linear; if two inches, 200 times; and so on.

To measure the size of objects, they may be placed directly on the micrometer; but as this is at all times inconvenient, whilst the object and micrometer, from their not being in the same plane, cannot, under high powers, both be brought into focus at once, it is better to use an eyemicrometer. Many ingenious inventions of this kind are



Spaces equal to 1-1000th of an inch magnified 250 diameters linear.



Five ruled spaces in an eyemicrometer, corresponding to one of those above, and consequently equal to the 1-5000th of an inch.

to be procured. The most simple is a ruled micrometer placed in the focus of the upper glass of the eyepiece. With this we observe how many divisions of the eye-micrometer correspond with one of those magnified by the microscope, always making our observation in the centre of the field, where the aberration of sphericity is least. On the latter being removed and replaced by an object, it becomes a matter of mere calculation to determine its size. Thus, supposing each of the upper spaces in Fig. 23 to represent the 1-1000th of an inch magnified 250 diameters linear,

and five of the lower spaces, as seen in an eye-micrometer, to correspond with one of these—it follows that each of these latter must measure 1-5000th of an inch. Oberhaeuser has made beautifully ruled eye-micrometers, for

the model recommended, which those who wish to make measurements would do well to procure.

If it be not in your power to estimate the magnifying power for yourself, the optician will give you a table, setting forth the various degrees of enlargement possessed by the lenses, and different eye-pieces, with the tube up or down. This table should always be referred to during the description of objects, and the amount of magnifying power invariably stated.

The art of demonstrating under the microscope is only to be acquired by long practice, and, like everything requiring practical skill, cannot be learnt from books or systematic lectures. I can only, therefore, give you very general directions on this head.

All that is necessary in examining fluid substances, is to place a drop in the centre of a slip of glass, and letting a smaller and thinner piece of glass fall gently upon it, so as to exclude air bubbles, place it upon the stage under the objective. In this way the fluid substance will be diffused equally over a flat surface, and evaporation prevented, which would dim the objective. The illumination must now be carefully arranged, and the focus obtained, first by means of the coarse, and then by means of the fine, adjustment. It will save much time, in examining structures, to employ always, at one sitting, the same slips of glass, as it is easier to clean these with a towel, after dipping them in water, than to be perpetually shifting the coarse adjustment.

The action of water, acetic acid, and of other re-agents, on the particles contained in a fluid, may be observed by mixing with it a drop of the re-agent before covering with the upper glass; or if this be already done, the drop of re-agent may be placed at the edge of the upper glass;

when it will be diffused through the fluid under examination by imbibition.

The mode of demonstrating solid substances will vary according as they are soft or hard, cellular or fibrous, etc. etc. The structure of a soft tissue, such as the kidney, skin, cartilage, etc., is determined by making very minute, thin, and transparent slices of it in various directions, by means of a sharp knife or razor. These sections should be laid upon a slip of glass, then covered over, and slightly pressed flat, by means of an upper one. The addition of a drop of water renders the parts more clear, and facilitates the examination, although it should never be forgotten that most cell-structures are thereby enlarged or altered in shape from endosmosis. Acid and other re-agents may be applied in like manner. The double-bladed knife of Valentin will enable you to obtain large, thin, and equable sections of such tissues, and permit you to see the manner in which the various elements they contain are arranged with regard to each other. . Harder tissues, such as wood, horn, indurated cuticle, etc., may be examined by small thin sections, made in the same way. Very dense tissues, such as bone, teeth, shell, etc., require to be cut into thin sections, and afterwards ground down to the necessary thinness. Preparations of this kind are now manufactured on a large scale, and may be obtained at a trifling cost. A cellular parenchymatous structure, such as the liver, may be examined by crushing a minute portion between two glasses. If it be membranous, as the cuticle of plants, epithelial layers, etc., the membrane should be carefully laid flat upon the lower glass, and covered with an upper one. A fibrous structure, such as the areolar, elastic, muscular, and nervous tissues, must be separated by means of needles, and then spread out into a thin layer before examination, with or without water, etc.

The commencing observer should not be discouraged by the difficulties he will have to encounter in dissecting and displaying many tissues. He must remember that the figures he sees published in books are generally either fortunate or very carefully prepared specimens. Practice will soon enable him to obtain the necessary dexterity, and to convince himself of the importance of this mode of inquiry. He should early learn to draw the various objects he sees, before and after the action of re-agents, not only because such copies constitute the best notes he can keep, but because drawing necessitates a more careful and accurate examination of the objects themselves. A note-book and a pencil for the purpose should be the invariable accompaniments of every microscope.

## How to observe with a Microscope.

The art of observation is at all times difficult, but is especially so with a microscope, which presents us with forms and structures concerning which we had no previous idea. Rigid and exact observation, therefore, should be methodically cultivated from the first, in order to avoid those errors into which the tyro, when using a microscope, is particularly liable to fall. Thus, you should carefully examine the physical properties of the particles and ultimate structures you may see, and not hastily conclude that you have under observation so-called pus, tubercle, or cancercorpuscles, because they were obtained from what was, a priori, believed to be pus, tubercle, or cancer. Nothing has been more clearly demonstrated by the progress of histology, than the fact, that the naked sight has confounded different structures together, from a similarity of external appearance, and that the greatest caution is required at all times, but especially by learners, in forming opinions as to the nature of different tissues

The physical characters which distinguish microscopic objects consist of—1st, Shape; 2d, Colour; 3d, Edge or border; 4th, Size; 5th, Transparency; 6th, Surface; 7th, Contents; and 8th, Effects of re-agents. These we may notice in succession.

- 1. Shape.—Accurate observation of the shape of bodies is very necessary, as many of these are distinguished by this physical property. Thus the human blood globules, presenting a biconcave round disk, are in this respect different from the oval corpuscles of the camelidæ, of birds, reptiles, and fishes. The distinction between circular and globular is very necessary to be attended to. Human blood corpuscles are circular and flat, but they become globular on the addition of water. Minute structures seen under the microscope may also be likened to the shape of well-known objects, such as that of a pear, balloon, kidney, heart, etc. etc.
- 2. Colour.—The colour of structures varies greatly, and often differs, under the microscope, from what was previously conceived regarding them. Thus the coloured corpuscles of the blood, though commonly called red, are in point of fact yellow. Many objects present different colours, according to the mode of illumination—that is, as the light is reflected from, or transmitted through their substance, as in the case of certain scales of insects, feathers of birds, etc. Colour is often produced, modified, or lost, by re-agents, as when iodine comes in contact with starch corpuscles, when nitric acid is added to the granules of chlorophyle, or chlorine water affects the pigment cells of the choroid, and so on.
- 3. Edge or Border.—The edge or border may present peculiarities which are worthy of notice. Thus it may be dark and abrupt on the field of the microscope, or so fine

as to be scarcely visible. It may be smooth, irregular, serrated, beaded, etc. etc.

- 4. Size.—The size of the minute bodies, fibres, or tubes which are found in the various textures of animals can only be determined with exactitude by actual measurement, in the manner formerly described. It will be observed, for the most part, that these minute structures vary in diameter, so that when their medium size cannot be determined, the variations in size from the smaller to the larger should be stated. Human blood-globules in a state of health have a pretty general medium size, and these may consequently be taken as a standard with advantage, and bodies may be described as being two, three, or more times larger than this structure.
- 5. Transparency.—This physical property varies greatly in the ultimate elements of numerous textures. Some corpuscles are quite diaphanous, others are more or less opaque. The opacity may depend upon corrugation or irregularities on the external surface, or upon contents of different kinds. Some bodies are so opaque as to prevent the transmission of the rays of light, when they look black by transmitted light, although they be white, seen by reflected light. Others, such as fatty particles and oil globules, refract the rays of light strongly, and present a peculiar luminous appearance.
- 6. Surface. Many textures, especially laminated ones, present a different structure on the surface from that which exists below. If, then, in the demonstration, these have not been separated, the focal point must be changed by means of the fine adjustment. In this way the capillaries in the web of the frog's foot may be seen to be covered with an epidermic layer, and the cuticle of certain minute fungi or infusoria to possess peculiar markings. Not unfrequently

the fracture of such structures enables us, on examining the broken edge, to distinguish the difference in structure between the surface and the deeper layers of the tissue under examination.

- 7. Contents.—The contents of those structures, which consist of envelopes, as cells, or of various kinds of tubes, are very important. These may consist of included cells or nuclei, granules of different kinds, pigment matter, or crystals. Occasionally their contents present definite moving currents, as in the cells of some vegetables, or trembling rotatory molecular movements, as in the ordinary globules of saliva in the mouth.
- 8. Effects of Re-agents.—These are most important in determining the structure and chemical composition of numerous tissues. Indeed, in the same manner that the anatomist with his knife separates the various layers of a texture he is examining, so the histologist, by the use of re-agents, determines the exact nature and composition of the minute bodies that fall under his inspection. Thus, water generally causes cell formations to swell out from endosmosis; whilst syrup, gum-water, and concentrated saline solutions, cause them to collapse from exosmosis. Acetic acid possesses the valuable property of dissolving coagulated albumen, and, in consequence, renders the whole class of albuminous tissues more transparent. Thus it operates on cell walls, causing them either to dissolve or become so thin as to display their contents more clearly. Æther, on the other hand, and the alkalies, operate on the fatty compounds, causing their solution and disappearance. mineral acids dissolve most of the mineral constituents that are met with, so that in this way we are enabled to tell with tolerable certainty, at all events the group of chemical compounds to which any particular structure may be referred.

## LECTURE V.

# PRINCIPAL APPLICATIONS OF THE MICROSCOPE TO DIAGNOSIS.

A perfect application of the microscope, for the purposes of diagnosis, can only be arrived at by obtaining, in the first instance, a complete knowledge of the tissues of plants and animals, both in their healthy and diseased conditions. The medical practitioner may be called upon to distinguish, not only the various structures which enter into every species of food, every kind of animal texture and fluid, and every form of morbid product, but he will frequently have to judge of these when more or less disintegrated, changed, or otherwise affected by the processes of mastication, digestion, expectoration, ulceration, putrefaction, maceration, etc. etc. In this place, however, I propose merely calling your attention to those points which are more likely to fall under your notice at the bed-side. No doubt, the practical applications of the microscope are daily extending, and whilst there are many points which may be said to be scarcely investigated, those which have been most so require to be further studied. At the same time, a careful and persevering examination of the morphological elements found in the various excreta of the body, as modified by different diseases, or by constitution and diet, cannot but prove of great importance in the present state of practical medicine. Hence, besides shortly discussing what is known, I shall especially indicate what are those subjects which may be elucidated by such of you whose previous histological observations qualify them for the task.

## SALIVA.

The readiest way of examining the saliva is to collect a drop of that fluid at the extremity of the tongue, and let it fall on the centre of a slip of glass. It should be allowed to remain quiescent for a minute or so, until most of the bubbles of air have collected in a mass on the surface. This should then be gently scraped off or placed aside with a needle, and the subjacent fluid covered with a thin glass. There will now be observed, with a magnifying power of 250 diameters linear—1st, The salivary corpuscles; 2d, Epithelial scales of the mouth; 3d, Molecules and granules.

1. The salivary corpuscles are colourless spherical bodies, with smooth margins, varying in size from the 1-3000th to the 1-1500th of an inch in diameter. They contain a round nucleus, varying in size, but generally occupying a third of the cell; and between this nucleus and the cell-wall are numerous molecules and granules, which communicate to the entire corpuscle a finely molecular aspect. The addition of water causes these bodies to swell out and enlarge from



Fig. 24.

endosmosis; while acetic acid somewhat dissolves the cellwall, and it becomes more transparent; while the nucleus

Fig. 24. Salivary corpuscles, epithelial scales, with molecules and granules, as seen in a drop of saliva.

250 diam.


appears more distinct as a single, double, or tripartite body. Both water and acetic acid also produce coagulation of the albuminous matter contained in the fluid of the saliva, which assumes the form of molecular fibres, in which the corpuscles and epithelial scales become entangled, and present to the naked eye a white film.

- 2. The Epithelial scales found in the saliva are derived from the mouth, and consist of flat plates, variously shaped, but generally presenting an oblong or squarish form, more or less curled up at the sides. Not unfrequently these have five or six sides, and are assembled together in groups, with their edges adherent. In size they vary from the 1-800th to the 1-500th of an inch in length. Embedded in their substance is a round or oval nucleus, together with numerous molecules and granules. Water produces no change in these bodies; but acetic acid renders the scale more transparent, and the nucleus to appear more distinct, with a darker edge.
- 3. Associated with the salivary corpuscles and epithelial scales are several molecules and granules, which vary in number in different people, and at various times of the day.


There may also be occasionally found in the saliva various foreign substances derived from the food,—such as granular debris of different kinds, starch globules or vegetable cells, muscular fasciculi, portions of areolar tissue, tendon, or spiral filaments, etc.—derived from pieces of texture which have adhered to the teeth during mastication.

The saliva may present various alterations, dependent on disease of the mucous membrane of the mouth and tongue. This, when ulcerated, causes an increase in the molecular and granular matter. Many of the epithelial scales also lose their transparent character and become opaque, from an augmentation of granular matter in their substance. Not unfrequently, under such circumstances, they give rise to confervoid growths, which mainly spring up in the debris

collected in the mouth, either on the surface of ulcers, in the sordes which collect on the teeth, gums, and tongue of individuals labouring under fever, or even in the inspissated mucus of persons who sleep for a considerable time with the



mouth open (Fig. 25). In infants, the tongue and cavity of the mouth are not unfrequently covered with a yellowish



floculent matter, constituting the disease named muguet by the French, in which sporules and confervoid filaments, in a high state of development, may be detected in considerable numbers (Fig 26).

In epithelial cancroid of the tongue, the epithelial scales exhibit a great tendency to split up and form fibres, and may frequently be found on the surface of the ulcer

presenting the form here figured (Fig. 27).

Occasionally they exhibit a tendency to form concentric

Fig. 25. Minute confervoid filaments springing from an altered epithelial scale, scraped from the surface of a cancroid ulcer of the tongue (*Leptothrix buccalis*).

Fig. 26. Confervoid filaments and sporules, in the exudation on the mouth and gums, constituting *Muguet* in infants.

Fig. 27. Fringe-like epithelium, from surface of an ulcer on the tongue.  $250\,di$ .

circles, or cell-nests, as they have been called, an appearance very common in the substance of the morbid growth, but also occasionally seen in the nodules which are from time to time separated and found in the saliva.




Fig. 28.

An histological examination of the saliva, of the fur and load of the tongue, in the great majority of diseases, is still a desideratum.

#### MILK.

On examining a drop of milk\* we observe a number of bodies rolling in a clear fluid. These bodies, in healthy milk, are perfectly spherical, with dark margins, smooth and abrupt on the field of the microscope, with a clear trans-

\* The mode of examining all fluids is the same, and is described p. 99; it need not be repeated.

Fig. 28. Concentric laminæ of epithelial scales, with epithelial cells and fragments of muscular fasciculi, from a cancroid ulcer of the tongue. 250 di.

parent centre, which strongly refracts light. In size they vary from a point scarcely measurable, up to, in different specimens, the 1-4000th or 1-3000th of an inch in diameter. In excess of ether they are dissolved or disappear; but if this re-agent be in small quantity, exosmosis takes place, and the field of the microscope is covered with loose globules of oil, of various forms. Water causes the milk globules to swell out, but very slightly. Acetic acid coagulates the caseous fluid in which they swim, and causes the globules to be aggregated together in masses. Several of the globules also exhibit, under the action of this re-agent, a certain flaccidity, and readily run into one another under pressure.

These globules consist of an albuminous delicate envelope, enclosing a drop of oil or butter. The membrane keeps them separate, so long as it is intact; but, dissolved by means of acetic acid, or ruptured by heat or mechanical violence (as in the churn), the butter is readily separated and collected. Cream is composed of the larger of these globules, which, owing to their light specific gravity, float on the surface of milk when allowed to repose.

The richness of milk is determined by the quantity of these globules. An examination of cow's and human milk will at once shew that the former contains a larger number than the latter. In all efforts, however, to determine the relative value of milk by microscopic examination, great care must be taken that the drop of fluid examined should be of the same bulk, that the same upper glass should be used in every case, and that it should be applied and pressed down with the same force. It is very difficult at all times strictly to fulfil these conditions, for not only is great skill in manipulation required, but an intimate acquaintance with the appearance of milk as seen under the microscope, is necessary, before any confidence can be placed in this mode of testing the quality of different specimens of the fluid. At the same time, the difference in the amount of oily consti-

tuents between the milk of the cow, ass, and human female, may in this way be easily determined.

In the same manner the various adulterations of milk are at once determined. Water, of course, separates the globules more and more from each other according to its amount. Flour will exhibit the large starch corpuscles, which are changed blue by the action of iodine. Chalk shews numerous irregular mineral particles, which are soluble in the mineral acids; and broken-down brain will be distinguished by large oil globules, mingled with fragments of fine nervetubes. Milk, when acid, exhibits the same character that it does under the action of acetic acid.

Healthy and fresh milk is indicated by a certain uniformity in the size of the globules; by their perfectly globular form; by their rolling freely over each other, and not collecting together in masses (Fig. 29). When the latter circumstance occurs, it is a sign of acidity.





The milk first secreted after parturition is called the colostrum. It is yellow in colour, and may be seen under the microscope to contain globules more variable in size, mingled with a greater or less number of compound granular bodies (Fig. 30.) These latter ought to disappear in the human female on the fifth or sixth day after parturition, but occasionally they remain, when the milk must be considered

Fig. 29. Globules of cow's milk.

Fig. 30. Colostrum of the human female, containing milk globules greatly varying in size, with granular corpuscles. 250 diam.


as unhealthy. In some cases I have seen them abundant so late as six weeks after the birth of the infant.

On some occasions, milk may be mixed with pus and blood, which are readily detected by the characters distinctive of each. Dr. Peddie has pointed out that milk can be squeezed from the mamma during the early months of pregnancy. Under such circumstances, it constitutes a most important sign of the pregnant state, especially of a first pregnancy; for although the secretion at this time has seldom the external appearance of milk, but is serous-looking, and often very viscid and syrupy, still, if examined with the microscope, the characteristic milk globules will at once appear. See his valuable paper, "Monthly Journal of Medical Science," August 1848.

#### THE BLOOD.

On examining a drop of blood drawn from the extremity of the finger by pricking it, there will be seen a multitude of yellow, round, bi-concave discs, rolling in the field of the microscope, which soon exhibit a tendency to turn upon their edge, and arrange themselves in rolls, like rouleaux of coins. These rouleaux, by crossing one another, dispose themselves in a kind of net-work, between which may be seen a few colourless spherical corpuscles, having a molecular surface, and a few granules. The coloured bloodcorpuscles vary in size from the  $\frac{1}{5000}$ th to the  $\frac{1}{3000}$ th of an inch in diameter, their average size being about the 4000 th of an inch—according to Gulliver, 3200 th of an inch. Owing to their bi-concave form, they present a bright external rim with a central shadowed spot, or a bright centre and a dark edge, according to the focal point in which they are viewed (Fig. 31). If the blood be exposed to the air a little time before examination, or if it be obtained by venesection, the edges of the corpuscles may often be observed to have lost their smooth outline, and to

have become irregular, notched, serrated, beaded, etc. (Fig. 32). Long maceration in serum, or other circumstances,



frequently cause them to diminish in bulk half their natural size, and present a perfectly spherical coloured body. On the addition of water, the blood discs become spherical, and lose their colour. On adding syrup, they become flaccid and irregular. Strong acetic acid dissolves them rapidly, and very weak acetic acid does so slowly, or diminishes their bulk by one-half.

The colourless corpuscles of the blood are spherical in form, and vary in size from the  $\frac{1}{2500}$  to the  $\frac{1}{2000}$  of an inch in diameter. Their surface presents a molecular or dotted aspect, which almost disappears on the addition of water, when they swell out by endosmosis. Acetic acid renders the external cell-wall very transparent, and brings the nucleus into view, consisting of one, two, or three round granules (Fig. 38.)

The examination of the blood by the microscope enables us to determine certain pathological conditions of that fluid, which, though few in number, are by no means unimportant.

In several diseases, the blood presents unusual spissi-

Fig. 31. Blood-corpuscles, drawn from the extremity of the finger. On the left of the figure they are isolated, some flat and on edge, some having a dark and others a light centre, according to the focal point in which they are viewed. On the right of the figure several rolls have formed. Two colourless corpuscles and a few granules are also visible.

Fig. 32. Blood-corpuscles altered in shape from exosmosis.

250 diam.

tude, depending on excesss of fibrin. In this condition the coloured blood-corpuscles easily lose under pressure their rounded margin, and assume a caudate or flask-like shape. They do not present their usual tendency to accumulate in rolls, but aggregate themselves together in irregular masses, as represented Fig. 33.





Fig. 33.

Fig. 34.

In certain internal hæmorrhages the blood-corpuscles break down, or become partially dissolved, when the external envelope is seen very transparent, the shadowed spots disappear, and there is found in their interior one or more granules. The liquor sanguinis also contains an unusual number of granules (Fig. 34). The same change is occasionally observable in the blood extravasated below the skin in scurvy or purpura hæmorrhagica.

In a woman who died of cholera, Dr. James M. Cowan observed a remarkable alteration in the blood, which he was so good as to show me. It consisted in the coloured corpuscles being paler than usual, and the colourless ones normal, but mingled with these were others varying in shape and size. They were generally circular, but some were oval, and a few caudate. They had a well-defined external smooth border, having one or two bright refracting granules, generally situated in the external membrane, and occasionally projecting from it. When seen edgeways,

Fig. 33. Blood-corpuscles altered in form, and aggregated together, in thick-ened blood.

Fig. 34. Altered blood corpuscles in the fluid of an hæmatocele. 250 diam.

they were flattened, and existed in the proportion of one to seven of the coloured corpuscles. Their long diameter varied from the  $\frac{1}{1200}$ th to the  $\frac{1}{2000}$ th of an inch, and their transverse diameter from the  $\frac{1}{2000}$ th to the  $\frac{1}{4000}$ th of an inch (Fig. 35). The addition of acetic acid caused them to swell out, dissolved their external wall, and liberated the granules. Aqua potassæ rendered the whole structure paler, and a solution of muriate of soda rendered them more distinct, and of smaller size.\*




Fig. 35.

Fig. 36.

We have seen that, in a healthy condition, the blood possesses very few colourless corpuscles, although sometimes




Fig. 37.

Fig. 38.

they are slightly increased in number, above the normal \* See Dr. Cowan's case.—Monthly Journal of Medical Science. March 1854.

250 diam.

Fig. 35. Appearance of blood once observed in a case of cholera.—(After Cowan.)


Fig. 36. Colourless corpuscles slightly increased in number.

Fig. 37. Appearance of a drop of blood, in Leucocythemia.


Fig. 38. The same, after the addition of acetic acid.

standard, as in Fig. 36, which may occur in a variety of dropsical and cancerous affections. But there is a certain state of that fluid I was the first to describe in 1845, and have since called "Leucocythemia," or white-cell blood, in which they are very numerous, generally associated with enlargement of the spleen or other lymphatic glands. The blood then presents the characters represented in the accompanying figures. (See Figs 37, 38.)

Occasionally, when increased in number, they are of



smaller size than those represented, Fig. 37, as in a case which entered the Infirmary with cancer of the thyroid body. (Figs. 39, 40.)



More rarely the nuclei of these corpuscles are found

- Fig. 39. Colourless corpuscles increased in number, and of small size.
- Fig. 40. The same, after the addition of acetic acid.
- Fig. 41. Numerous naked nuclei of the colourless corpuscles in the blood.
- Fig. 42. The same, after the addition of acetic acid.

250 diam.


naked in the blood, as in cases recorded by Virchow and myself.\* (Figs. 41, 42.)

It has been affirmed that the colour and number of the corpuscles of the blood undergo a change in plethora, fever, jaundice, dropsies, cholera, etc., but exact observations are wanted to confirm the statement. I have never been able to satisfy myself that any such changes were observable in these diseases by means of the microscope. In chlorosis the number of the blood-globules is undoubtedly diminished; but this is determined by the size of the clot, rather than by microscopic demonstration.

Occasionally the serum of the blood presents a lactescent appearance; and, on being allowed to remain at rest some hours, a white creamy pellicle forms on the surface. This consists of very minute particles of oil, which resemble the smaller molecules found in milk and in the chyle.

#### Pus.

Normal or good pus, when examined under a microscope, is found to consist of numerous corpuscles, floating in a clear fluid, the *liquor puris*. The corpuscles are globular



in form, having a smooth margin, and finely granular surface. They vary in size from the  $\frac{1}{2000}$ th to the  $\frac{1}{1200}$ th of an inch in diameter. There may be generally observed

\* See the author's work "On Leucocythemia, in relation to the Physiology and Pathology of the Lymphatic Glandular System." Edinburgh, 1852.

Fig. 43. Pus corpuscles, as seen in healthy pus.

Fig. 44. The same, after the addition of acetic acid.

in some of them a round or oval nucleus, which is very distinct on the addition of water, when also the entire corpuscle becomes distended from endosmosis, and its granular surface is more or less diminished. On the addition of strong acetic acid the cell-wall is dissolved, and the nuclei liberated in the form of two, three, four, or rarely five granules, each having a central shadowed spot. If, however, the re-agent be weak, the cell-wall is only rendered very transparent and diaphanous, through which the divided nucleus is very visible (Fig. 44).

Occasionally these bodies are seen surrounded by another fine membrane, as in Fig. 45. At other times they are not perfectly globular, presenting a more or less irregular margin, associated with numerous molecules and granules.



This occurs in what is called scrofulous pus, and various kinds of unhealthy discharges from wounds and granulating surfaces. (Fig. 46.) In gangrenous and ichorous sores, a few of these irregular corpuscles are associated, not only with a multitude of molecules and granules, but with transformed and broken-down blood globules, the debris of the involved tissues, etc. etc.

## SPUTUM.

A microscopic examination of the sputum demands a most extensive knowledge of both animal and vegetable structure. I have found in it—1st, All the tissues which enter into the composition of the lung, such as filamentous tissue, young and old epithelial cells, blood-corpuscles, etc.

Fig. 45. Pus corpuscles, surrounded by a delicate cell-wall.

Fig. 46. Irregular-shaped pus corpuscles, in scrofulous pus. 250 diam.

2d, Mucus from the œsophagus, fauces, or mouth. Morbid growths, such as pus, pyoid, and granular cells; tubercle corpuscles, granules, and amorphous molecular matter; pigmentary deposits of various forms, and parasitic vegetations, which are occasionally found in the lining membrane of tubercular cavities. 4th, All the elements that enter into the composition of the food, whether animal or vegetable, which hang about the mouth or teeth, and which are often mingled with the sputum, such as pieces of bone or cartilage, muscular fasciculi, portions of esculent vegetables, as turnips, carrots, cabbages, etc.; or of grain, as barley, tapioca, sago, etc.; or of bread and cakes; or of fruit, as grapes, apples, oranges, etc. All these substances render a microscopic examination of expectorated matters anything but easy to the student.

To examine sputum, it should be thrown into water, when, on account of the air it contains, it will generally float on the surface; while the more dense portions, such as masses of crude tubercle or cretaceous concretions fall to the bottom. It should be then teased, or broken up with a rod, when the various elements and particles it contains will gradually disengage themselves, and may be separated from the mass without difficulty. Nothing is more common, on examining portions of sputum with a microscope, than to











observe the various aggregations of molecular and granular matter here figured.

Fig. 47. Mass consisting of minute molecules, frequently seen in disintegrated tubercle.


Figs. 48 and 49. Masses composed of molecules and oily granules varying in size and mode of aggregation.

Fig. 50. Mass partly composed of the debris of a fibrous structure.

Fig. 51. Mass composed of tubercle corpuscles.

250 diam.

Occasionally little masses of a cheesy substance, and yellowish colour, may be found entangled in the purulent mucus, or collected at the bottom of the vessel. These, when examined, present a number of irregular-shaped bodies, approaching a round, oval, or triangular form, varying in their longest diameter from the 1-4000th to 1-2000th of an inch. These bodies contain from one to seven granules, are unaffected by water, but are rendered very transparent by



acetic acid. They are what have been called tubercle corpuscles. They are frequently mingled with a multitude of molecules and granules, which are more numerous in proportion to the softness of the tubercle (Figs. 51 and 52.)


Sometimes indurated or gritty little masses are brought up with the sputum, which are derived from the cretaceous or calcareous transformation of chronic tubercle in the lungs. They consist of irregular masses of phosphate of lime, combined with more or less animal matter. On squeezing such as are friable between glasses, and examining their structure, they frequently may be seen to contain the elements represented in Fig. 53.

Fig. 52. Tubercle corpuscles and granules, from a softened tubercular mass in the lungs, frequently found in the sputum.

Fig. 53. Fragments of phosphate of lime, crystals of cholesterine, and tubercle corpuscles, from a cretaceous mass in the lungs, occasionally found in the sputum.

250 diam.

Sputum frequently presents a fibrillated appearance, which is common to all mucous discharges. This is caused by the deposition in viscid mucus of molecules, which assume a linear arrangement. This deposition is increased by the addition of water and acetic acid, so that they consist of albumen. These fine molecular fibres (see Figs. 59, 70, 71, and 74) must be separated from the areolar and elastic tissue of the lung, which is not unfrequently found in sputum, indicating ulceration or sloughing of the pulmonary texture. Shroeder van der Kolk has lately stated, that these fragments may be found in the sputum before the




physical signs of ulceration of the lung, as determined by auscultation, are well characterised. This fact I have confirmed, and believe it to be one of great diagnostic importance.

In acute pueumonia and what is called fibrinous bronchitis the sputum frequently contains fibrinous casts of the minute bronchi, which present a branched mould of the

Fig. 54. Fragment of elastic tissue of the lung, in phthisical sputum.

Fig. 55. Fragment of areolar and elastic tissue, still exhibiting the form of air cells from phthisical sputum. a, Another fragment.—(Van der Kolk) 250 di.

tubes. These casts may be readily separated in water, as previously described; and when examined with the micros-



cope, are found to consist of molecular fibres, in which pyoid and pus corpuscles are infiltrated. (Fig. 57.)

The inspissated sputum, so commonly expectorated in the morning, is derived from the fauces. It often presents a dirty green or brownish colour, passing into black. When examined with a microscope, it may be seen to consist of epithelial cells, more or less compressed together, and varying in size from the 1-2000th to the 1-800th of an inch in diameter. The smaller ones are round, and closely resemble pus corpuscles; the larger ones are round or oval, with a distinct nucleus. In the dark-coloured portions of this sputum, the cells contain numerous granules and molecules, several of which are black and quite opaque. This black matter consists of carbon, and is unaffected by re-agents. The addition of acetic acid causes coagulation of the mucus

Fig. 56. Fibrinous coagula in sputum, exhibiting moulds of the bronchi. Natural size.—(After Peacock.)

Fig. 57. Fibres, with corpuscles, in a fibrinous coagulum from a bronchus.

in which the cells are embedded; and whilst it produces little change in the older cells, it dissolves, or renders tran-



Fig. 58.




Fig. 59

sparent, the walls of such as are young, displaying a round, oval, or divided nucleus, as seen in the figure. (Fig. 53.)

In the black phthisis of colliers the sputum is ink-black, and more or less tenacious. On examination with a microscope, the cells are seen to be loaded with carbonaceous




Fig. 60.

pigment. Several of these cells are perfectly opaque, whilst others are almost colourless; and between the two extremes there is every kind of gradation as to intensity of blackness. This black pigment is unaffected by the strongest re-agents,

Fig. 58. Epithelial cells, embedded in mucus, expectorated from the fauces. Some are seen to contain black pigment; others resemble pus corpuscles.

Fig. 59. Another portion of expectorated mucus from the fauces, acted on by acetic acid, showing fibrillation and the changes in the young cells.

Fig. 60. Cells in the black sputum of the collier, the pigment in which is persistent under the action of every known chemical agent. 250 diam.

nitro-muriatic acid, chlorine, and even the blow-pipe, failing to decompose it. It is, therefore, pure carbon, and differs from the pigment contained in cells of similar appearance in melanotic tumours, as in these latter the re-agents just mentioned at once destroy the colour.

#### VOMITED MATTERS.

The matters rendered by vomiting have not been made so frequent an object of microscopical observation as is necessary, with a view to diagnosis. In organic diseases of the organ, nothing has been ascertained on this head. In other cases, it almost always happens, that the matters rendered consist—1st, Of the food and drink, in various stages of decomposition and disintegration; 2d, Of alterations in the epithelial lining membrane of the stomach, œsophagus or pharynx, mingled with more or less mucus; 3d, Of certain new formations, which are produced in the fluids of the stomach.

1. It would constitute a very interesting series of observations, to determine, with the aid of the microscope, the structural changes which various articles of food undergo during the process of digestion in the stomach. This has not yet been done with accuracy, although there can be little doubt that compound tissues become disintegrated in the inverse order to that in which they are produced—that is to say, fibres become separated, embedded cells become loose, and, when aggregated together, their cohesion is destroyed. The cell-walls then dissolve, the nucleus still resisting the solvent process for some time; but at length the whole is resolved into a molecular and granular mass, which in its turn becomes fluid. Such, however, is the different soluble properties of various edible substances, that, in a time sufficient for the perfect solution of some, others are scarcely affected. It may readily be conceived, that the transitions which these substances undergo, may

occasionally render their detection difficult; and such is really the case. Starch corpuscles, for instance, break down



Fig. 61.

into rounded granules or molecules, and are very liable to puzzle an inexperienced observer. Tincture of iodine, from its peculiar reaction on these bodies, will always enable us to recognise them.

2. The various epithelial cells which line the passages leading to the stomach, as well as the structures peculiar






Fig. 63.

to that organ itself, may be found in the vomited matters —of course mingled with the debris of edible substances. They also may have undergone various changes in appear-

Fig. 61. Appearance of starch corpuscles after partial digestion in the stomach.

Fig. 62. Flake in the rice-water vomiting of a cholera patient, showing, a, large epithelial cells; b, milk globules, and coagulated caseine; c, torulæ; and d, half-digested epithelial scales, with liberated nuclei, more or less broken down.

Fig. 63. Structures observed in certain rice-water vomitings from a cholera patient, showing bodies which consist of the half-digested uredo in bread. 250 di. ance, from endosmosis, or even partial digestion. In cholera, the vomited matter consists principally of such altered epithelial cells or scales, many of which are derived from the fauces or esophagus.

3. The new formations which may be produced in the stomach are principally vegetable fungi—such as various kinds of torulæ (see Fig. 62, c), and especially one first discovered in vomited matters by Mr. Goodsir, and which he has called, on that account, Sarcina Ventriculi. It consists




Fig. 64.

of square particles, which apparently increase by fissiparous division in regular order, so that they present square bundles of four, sixteen, or a multiple of these. Although at first supposed to be peculiar to the stomach, I have frequently found them in the fœces; and in one case, in the urine. They were also found by Virchow, in an abscess of the lung.

In addition to the bodies alluded to, occasionally observed in vomited matters, they may contain various morbid products, such as blood, pus, and cancer-cells, colouring matter of the bile, etc.

#### Fœces.

The same difficulty attends the examination of the fœces as of the sputum; for there may be found in it,—lst, All the parts which compose the structure of the walls of the alimentary canal; 2d, All kinds of morbid products; and, 3d, All the elements which enter into the composition of food. The only difference is, that these last are generally more broken down or disintegrated.

Under certain circumstances, the diagnostic value attached to the examination of the fœces is greater than that of the sputum, or of vomited matters. For instance, when pus or blood globules are detected, we may infer that the more perfect these are, the nearer to the anus did they originate. In examples 4 and 6 (pp. 83, 84) I have shown how the detection of certain vegetable structures, used as food, were serviceable in diagnosis; but this subject merits more extensive attention than has hitherto been paid to it.

In typhus, and other putrid fevers, the stools contain masses of large crystals of phosphates or carbonates. In dysentery they are loaded with pus and blood; and the former may also be detected on the surface of feecal masses when the intestine is ulcerated. There may also be observed




Fig. 65.

numerous torulæ, and occasionally sarcinæ. In cholera the white stools consist of mucus, in which the debris of epithelial cells are entangled; and as the nuclei of these cells resist disintegration for a long time, these round or oval bodies generally exist in considerable numbers. (Fig. 65.)

In a disease very common in Edinburgh, especially in women, in which flakes of membranous matter are thrown off from the bowels in large quantities, these present a very similar appearance to the cholera flakes just noticed.

Among the indigestible articles connected with the food, it was observed, in the autumn of 1849, that curious-shaped bodies were detectable, both in the vomited matters and stools of cholera patients. These were supposed to be parasitic formations connected with the cause of cholera, but were pointed

Fig. 65. Structure of flakes in a rice-water stool, from a cholera patient.

out by Mr. Busk to be the *uredo-segitum*, occasionally found in bread. (Figs. 63 and 66.)



Fig. 66.

On one occasion, a dispensary patient brought to me a membranous mass, which had been evacuated by the bowels. It resembled a piece of boiled fine leather, of a greenishyellow colour, and fibrous structure. On microscopic ex-



Fig. 67.

Fig. 68.

amination, it was found to be made up of an inextricable mesh-work of confervoid growths, consisting of long tubes, with joints, and a few oval sporules, the former having a great tendency to break across. (Figs. 67 and 68.)

## UTERINE AND VAGINAL DISCHARGES.

The diagnostic indications to be derived from the microscopic examination of these discharges, has not been much investigated; but there are few subjects which hold out the promise of more useful results to the medical practitioner. It can only be prosecuted by the obstetric histologist, who,

Fig. 66. Portions of the uredo in bread, still further digested and disintegrated than is observable in the vomited matters. (Fig. 63.) Some torulæ are also present.

Fig. 67. Structure of confervoid mass passed from the bowels.

Fig. 68. The same, magnified 500 diameters linear, showing their vegetable nature. 250 diam.

on collecting the secretions poured out from the os uteri, or on the vaginal walls, by means of the speculum, should observe their structural peculiarities when quite fresh.\*

The menstrual discharge will be found to consist of young epithelial cells, old epithelial scales, and blood globules, the number of which last will be greater or less accord-




Fig. 69.

ing to the intensity of the colour. A leucorrheal discharge always consists of old epithelial scales, which may be more or less loaded with fat, combined with numerous young epithelial cells (round or oval), and pus corpuscles. (Fig. 69.)


The white gelatinous discharge, so frequently seen, with the speculum to be derived from the os uteri, consists of gelatinous mucus, in which round or oval young epithelial cells are mingled. The mucus is copiously deposited in a molecular form, on the addition of acetic acid or water, whilst the walls of the cells are rendered transparent, and an oval granular nucleus made visible. (Figs. 70 and 71.)

Not unfrequently leucorrheal and other discharges contain groups of blood-globules, the shapes of which are almost always more or less altered by endosmosis, on account of the

\* On this point Dr. Tyler Smith's work on Leucorrheea may be consulted with advantage.


Fig. 69. Corpuscles seen in a chronic leucorrheal discharge, consisting of,-1st, Large epithelial scales, from the vagina and cervix uteri. On the left of the figure some of these may be observed to have undergone the fatty degeneration. 2d, Numerous pus corpuscles; and, 3d, Blood globules, the external edges of which are more or less dentated from exosmosis. 250 diam.

viscid fluid mingled with them (see Fig. 69). Indeed, the variations observable in these discharges are dependent for



the most part on the excess of one or more of the elements just mentioned—namely, epithelial cells or scales, pus or blood corpuscles, and gelatinous mucus.

In addition to the fluid discharges poured out from the uterus and vagina, there are a variety of morbid growths connected with these organs, the diagnosis of which may be materially facilitated by microscopic examination. The sepa-



ration of fibrous, epithelial, and cancerous tumours and ulcers belong to this category, which must be conducted on the general principles referable to the diagnosis of morbid

Fig. 70. Structure of gelatinous mucus from the os uteri.

Fig. 71. The same, after the addition of acetic acid.

Figs. 72 and 73. Two specimens of cancerous juice squeezed from the uterus.

250 diam.

growths in general. I have had abundant opportunities of satisfying myself of the importance of this mode of proceeding, in cases where the substance, mucous surface, or cervix of the uterus has been more or less involved.

#### Mucus.

In all fluids secreted from a mucous membrane, many of which have been noticed, there may be found a gelatinous material, which has long been called mucus. It may vary in colour from a milk-white to a yellowish brown or even black tint, these variations being dependent on the cell structures or pigment it contains. By some it has been supposed that there are certain cell formations peculiar to mucus, which have been called "mucus corpuscles;" but it has always appeared to me that the various bodies found in this secretion are either different forms of epithelium, on the one hand, or pus cells on the other. Thus the round epithelial cells found in mucous crypts, or the bodies constituting permanent epithelium, when newly formed, before they have had time to flatten out, and perhaps more or less affected by endosmosis, are represented, Figs. 52 and 65. These are the mucous corpuscles of some writers. when exudation is poured out on a mucous surface, and is mingled in greater or less quantity with the gelatinous secretion, it presents a marked tendency to be transformed into pus corpuscles, and hence why all irritations of mucous surfaces are usually accompanied by purulent discharges. The pus corpuscles, under such circumstances, present all the characters formerly noticed as peculiar to these bodies (see Figs. 43 and 69).

Hence, properly speaking, there is no such body as a mucus corpuscle, the cells found in mucus being either epithelial or puscells, the number of which present communicates certain peculiarities to the discharge. Thus, as we have seen, the white gelatinous mucus discharged from the os uteri

contains the former, whilst the peculiar fluid characteristic of a gonorrhœa or catarrh, in either sex, abounds in the latter. The gelatinous substance, however, in which these bodies




Fig. 74.

are found (Mucin) is what is peculiar to the fluid secreted from mucous surfaces, containing, as it does, a large amount of albumen possessing a remarkable tendency to coagulate in the form of molecular fibres. (Fig. 74.) When recent, these are few in number, but on the addition of water or acetic acid they are precipitated in such numbers as to entangle the cell formations, and present a semi-opaque membranous structure. (Figs. 59, 70, and 71.)

The more healthy a mucous secretion, the more it abounds in this viscous albuminous matter, and the less are its cell elements. On the other hand, when altered by disease, the cell elements increase, and the viscosity diminishes.

## DROPSICAL FLUIDS.

The fluids obtained by puncture of dropsical swellings, may in some cases, when examined microscopically, present peculiarities worthy of notice. Thus, in the serum collected within the tunica vaginalis testis, numerous spermatozoa

Fig. 74. Viscid, greyish yellow sputa of pneumonia, treated with dilute acetic acid, containing fibrinous mucin, pus corpuscles, and epithelial cells containing fatty and pigment granules.—(After Wedl.) 300 diam.


may be found, constituting what has been called spermatocele. How these bodies find their way into this fluid is unknown, as no direct communication with the substance



of the testicle has ever been seen; neither does their occurrence seem to interfere in any way with the successful treatment of this kind of dropsy, by injections, as practised in hydrocele.

In the fluid of ascites, when removed from the body, there may usually be observed a few epithelial scales from the serous layer of the abdomen, which are more abundant in some cases than in others. Occasionally blood and pus corpuscles may be detected in greater or less quantity.

In ovarian dropsy, various products may be found in the



evacuated fluid, according to the nature of the contents of the cyst. Pus and blood corpuscles are common elements, but more commonly epithelial cells and scales, which occa-


Fig. 75. Spermatozoa as observed in the fluid of Spermatocele.

Fig. 76. Cells in fluid, removed from an ovarian dropsy.


Fig. 77. The same, after the addition of acetic acid.

Fig. 78. Group of similar cells, many-sided from compression. 250 diam.

sionally accumulate in the cysts of ovarian tumours. (Figs. 76, 77, 78.) At other times, masses of gelatinous or colloid matter are evacuated, which may present various appear-



ances, according to the time it has been secreted. Occasionally it exhibits a faintly fibrous structure, the large meshes of which are filled with transparent jelly, in which round or oval corpuscles, isolated or in groups, may be seen as represented in Figs. 79, 80.



Occasionally this substance is more or less opaque, and on examination, will be found to consist sometimes of fatty granules, either alone or combined with the flat characteristic crystals of cholesterine. (Figs. 81, 82.)

- Fig. 79. Group of oval corpuscles in the gelatinous matter of ovarian dropsy.
- Fig. 80. Round and oval corpuscles, with fibres in another gelatinous mass.
- Fig. 81. Fatty granules occasionally found in colloid masses, taken from an ovarian tumour.

Fig. 82. Fatty granules, and compound granular cells, with crystals of cholesterine, frequently found in similar masses. The exact angle of the crystals has not been well copied by the artist.

250 diam.


In the examn ation of dropsical fluids, also, there can be little doubt that further research will lead to very important results in diagnosis.

#### URINE.

Healthy human urine examined with the microscope, when recently passed, is absolutely structureless. Allowed to repose for twelve hours, there is no precipitate; occasionally a slight cloudy deposition may be observed, in which may be discovered a few epithelial scales from the bladder, a slight sediment of granular urate ammonia, or a few crystals of triple phosphate. In certain derangements of the constitution, however, various substances are found in the urine, which, in a diagnostic point of view, are highly important, and which we shall shortly notice in succession.


To examine the deposits found in urine, this fluid should be poured, in the first instance, into a tall glass jar; then decant the clear liquid, and put the lower turbid portion into a tall test tube, and again allow the deposit to form. this manner, the structural elements accumulate in the smallest possible compass, and a large number of them are brought into the field of the microscope at once. The quantity of any salt or deposit in the urine can never be ascertained by the microscope. But in the great majority of cases the appearances observed with that instrument, are sufficient in themselves to distinguish the nature of the various kinds of sediment met with, and these consequently are all that need be described in this place.

Uric Acid.—Uric acid crystals are almost always coloured, but the tint varies from a light fawn to a deep orange red. The general colour is yellow. They present a great variety of forms, the most common being rhomboidal. The lozengeshaped and square crystals, which are more rarely met with, isolated and in groups, are represented, Fig. 83. Not unfrequently they present adhering masses or flat scales with transverse or longitudinal markings, as seen, Fig. 84. Occa-



sionally they assume the form of truncated rounded columns, as represented, with other structures, Fig. 89.

Urate of Ammonia most commonly assumes a molecular and granular form, occurring in irregular aggregated amorphous masses. Fig. 86. This may be separated from a similar-looking deposit of phosphate of lime, by the action of dilute muriatic acid, which immediately dissolves the last-named salt, but acts slowly on urate of ammonia, setting free the uric acid. Sometimes, however, it occurs in spherical bodies of a bister brown colour, varying in size from the 1-5000th to the 1-2000th of an inch in diameter. The lat-



ter size rarely occurs. Occasionally they assume a stellate form, from needle-like or spicular prolongations coming off from the spherical body. I have seen both these forms associated, and the former so curiously aggregated together as to assume the appearance of an organic membrane, for

Fig. 83. Lozenge-shaped and rhomboidal crystals of uric acid.

Fig. 84. Aggregated and flat striated crystals of uric acid.

Fig. 85. Urate of ammonia, in a granular membranous form, and in rounded masses, with spicula. 250 diam.

which by some it was mistaken, until it was observed to dissolve under the action of dilute nitric acid. (Fig. 85.)

Triple Phosphate or Ammonio-Phosphate of Magnesia.—
These crystals are very commonly met with in urine, and are generally well defined, presenting the form of triangular prisms, sometimes truncated, at others having terminal facets. (Fig. 86.) If an excess of ammonia exist, or be added artificially, they present a star-like or foliaceous appearance, which, however, is seldom seen at the bed-side.

Most of the forms of urate of ammonia are represented,






Fig. 87.

Figs. 85 and 86, in the latter associated with the triple phosphate.

Oxalate of Lime most commonly appears in the form of octohedra, varying in size, the smaller aggregating together in masses. Once seen, these bodies are readily recognised. (Fig. 87.) Very rarely they present the form of dumb bells

or an oval body, the central transparent portion of which presents a dumbbell shape, while the shadowed dark portion fills up the concavities.

Cystine forms flat hexagonal plates, presenting on their surface marks of similar irregular crystals (Fig. 88.) Occasionally the centre is opaque, with




Fig. 88

radiations more or less numerous, passing towards the circumference.


Fig. 86. Triple phosphate, with various forms of urate of ammonia.

Fig. 87. Octohedral and dumb-bell shaped crystals of oxalate of lime.


Fig. 88. Flat and rosette-like crystals of cystine.

250 diam.

In addition to the various salts found in the urine, there may occasionally be found different organic products, such as blood and pus corpuscles, spermatozoa, vegetable fungi, exudation casts of the tubes, or epithelial scales from the bladder or mucous passages. Frequently one or more of these are found together, as in the following figure:—



Very rarely casts of the tubes, principally composed of oily granules, may be seen, or epithelial cells, more or less loaded with similar granules, several of which also float loose in the urine, as in the accompanying figure:—



Although these casts of the tubes were at one time con-

Fig. 89. Bodies observed in the urine of a scarlatina patient, 24 hours after being passed. a, Desquamated fragment of uriniferous tube. b, Exudation casts of uriniferous tubes. c, Amorphous urate of ammonia. d, Columnar crystals of uric acid. e, Blood corpuscles. f, Fus corpuscles. g, Torulæ and vegetable fungi, which had been formed since the urine was excreted.

Fig. 90. Cast of a uriniferous tube, principally composed of oil granules, with fatty epithelial cell and free oil granules, in urine of Bright's disease. 250 di.

founded together, they may now be separated into at least four distinct kinds, namely,—1st, Fibrinous or exudative; 2d, Desquamative; 3d, Fatty; and 4th, Waxy casts. The fibrinous and desquamative casts are most commonly found in urine in acute diseases of the kidneys, or at critical periods of severe inflammations, such as scarlatina, small pox, pneumonia, etc. The fatty and waxy casts are more indicative of chronic disease of the renal organs themselves. At the same time it should be understood that they may be more or less associated together.

Spermatozoa are occasionally found in the urine, but must not be considered as of any importance, unless accompanied by the peculiar symptoms of spermatorrhæa. (See Fig. 75.) The presence of torulæ in considerable quantity (Fig. 89, g) is indicative of the existence of sugar, which requires, however, for its confirmation, the application of chemical tests.

All the various appearances noticed are only diagnostic when accompanied by concomitant symptoms. Alone, they are not to be depended on; but, in combination with the history and accompanying phenomena, are capable of affording the greatest assistance in the detection of disease.

## CUTANEOUS ERUPTIONS AND ULCERS.

An examination of the various products thrown out upon the skin in the different forms of eruption, ulcer, and morbid growth, may in many cases be of high diagnostic value. Of these we shall speak separately.

1. Cutaneous Eruptions.—In the vesicular and pustular diseases, there may be observed below the epidermis all the stages of pus formation, commencing in exudation of the liquor sanguinis, gradual deposition of molecular and granular matter, and formation around them of cell-walls. The eruption produced artificially by tartar emetic ointment offers the best opportunity of examining the gradual forma-

tion of these bodies under the microscope. Pus taken from all kinds of eruptions and sores presents the same characters, there being no difference between the pus in impetigo, and that in variola. When a scab is formed, as in eczema or impetigo, a small portion of it broken down, mixed with water and examined under the microscope, presents an amorphous collection of granules, oil globules, and epithelial scales.

The squamous eruptions of the skin are three in number -namely, psoriasis, pityriasis, and ichthyosis. The dry incrustations which form on the surface in these diseases essentially consist of epidermic scales more or less aggregated together. They are very loose in pityriasis, and occa-




Fig. 91.

sionally mingled with debris of vegetable confervæ, similar to what grows on the mucous membrane of the mouth (Figs. 25 and 26). The scales are more aggregated together in psoriasis, and greatly condensed in ichthyosis; occasionally in the latter disease presenting the hardness and structure of horn.

The epidermic tumours of the skin assume the form of corns, callosities, condylomatous warts, and what has been called Veruca Achrocordon. They all consist, in like manner, of epidermic scales more or less condensed together; in the latter growth they surround a canal furnished with blood vessels. Sometimes they entirely assume a regular form, their interior being more or less hard, fibrous, and vascular,

-in short, a prolongation of the epidermis. At other times




Fig. 92.


they soften on their summits, and assume the structural peculiarities of the epithelial ulcer afterwards to be described.

The favus crust is composed of a capsule of epidermic scales, lined by a finely granular mass, from which millions of cryptogamic plants spring up and fructify. The presence of these parasitic vegetations constitutes the pathognomonic character of the disease.

In order to examine the natural position of these vegetations microscopically, it is necessary to make a thin section of the capsule completely through, embracing the outer layer of epidermis, amorphous mass, and light friable matter found in the centre. It will then be found, on pressing this slightly between glasses, and examining it with a magnifying power of 300 diameters, that the cylindrical tubes (thalli) spring from the sides of the capsule, proceed inwards, give off branches dichotomously, which, when fully developed, contain at their terminations (mycelia), a greater or less number of round or oval globules (sporidia). These tubes are from the

Fig. 92. The summit of a papilla from an epidermic growth, the result of a burn. Each papilla consisted externally of numerous epidermic scales distinctly nucleated, compressed together. Internally it was composed of fibrous vascular tissue. 250 diam.

1-6000th to 1-8000th of an inch in thickness, jointed at irregular intervals, and often containing very minute molecules in their interior. The longitudinal diameter of the sporules is generally from 1-4000th to 1-2000th, and the



transverse from 1-4000th to 1-6000th of an inch in diameter. I have seen some of these oval and round, twice the size of the others. The long diameter of the former measured 1-1500th of an inch. The mycelia and sporules agglomerated in masses are always more abundant and highly developed in the centre of the crust. The thalli, on the other hand, are most numerous near the external layer. There may frequently be seen swellings on the sides of the jointed tubes, which are apparently commencing ramifications.

On examining the hairs which pass through the favus crusts, it will often be found that they present their healthy structure. At other times, however, they evidently contain

Fig. 93. Branches of the Achorion Schoenleini, in an early stage of development, growing from a molecular matter, and mingled with epidermic scales, from a very minute favus crust.

Fig. 94. Fragments of the branches more highly developed, with numerous sporules and molecular matter, from the centre of an advanced favus crust.

300 diam.

long, jointed branches, similar to those in the crust, running in the long axis of the hair, which is exceedingly brittle. I have generally found these abundant in very chronic cases; and on adding water, the fluid may be seen running into these tubes by imbibition, leaving here and there bubbles of air, more or less long. There can be very little doubt, that the tubes and sporules after a time completely fill up the




Fig. 95.


hair follicle, and from thence enter the hair, causing atrophy of its bulb, and the baldness which follows the disease. The various steps of this process, however, I have been unable to follow, never having had an opportunity of observing favus in the dead scalp, and of making proper sections of the skin.

The skin is also attacked by certain animal parasites. Of these the pediculi, or lice, are too well known to need description. But we may shortly allude to the Acarus scabiei, and the Entozoon folliculorum.

The Acarus scabiei (see Fig. 96) has been proved by the researches of M. Bourguignon to be the undoubted cause of itch. The male is about a third smaller than the

Fig. 95. a, A light hair, containing branches of the Achorion Schoenleini—(magnified 300 diameters linear.) The wood-cutter has made the branches too beaded. b, A darker coloured hair, containing branches of the plant—(magnified 800 diameters linear).

female. He has suckers on two of his hind feet, and possesses on the abdominal surface genital organs, all of which characters are absent in the female. She, on the other hand, in addition to her size, and the negative marks alluded to, is



characterised by the three kinds of horny spines which are scattered over the back. The suckers, or ambulacria, are organs of locomotion; the mandibles enable it to cut the epidermis, and extract fluid from the tissues, which passes through a delicate esophagus, the internal termination of which is unknown, the body of the animal being apparently


Fig. 96. Female and male Acarus Scabies, magnified 100 diameters (after Bourguignon).

filled with an unorganised, very finely molecular pulp. A short delicate tube may also sometimes be observed at the anus—a supposed rectum. No respiratory apparatus can be discovered, although the creature may be seen to swallow minute bubbles of air, which pass down the œsophagus, and, like the nutritive juices, diffuse themselves through the interior. At all events, animal juice and air are both necessary to the life of the Acarus.

The male is more agile than the female, and may be observed to run about on the surface, plunging his head now and then into the skin to extract nourishment. But he never buries his body in holes or furrows as the female does, in order to deposit her eggs. In number the male is to the female as one to ten. He was first discovered by Eichstedt, subsequently by Lanquetin, and has been carefully studied, described, and figured by Bourguignon.

The habits of these insects account for the well known tegumentary irritation of persons affected with itch, when they get warm in bed or approach a fire. It appears that both males and females are active when the infected individual enjoys a certain elevated temperature. The males and young females then run about, and feed themselves by puncturing the skin and swallowing the juices and blood globules. It is then also that the mothers burrow into the epidermis, cutting with their mandibles, and pushing their bodies forward with the assistance of the spines on their backs which serve as points d'appui, and so form grooves which advance at about the rate of one-twelfth of an inch each night. The male, besides hunting for food, is very active in paying attention to his female friends, carefully avoiding, however, such as are about to become mothers. All these various operations sufficiently account for the itching on the surface of the skin, as well as the tingling and occasional stinging sensations often experienced.

The Entozoon Folliculorum inhabits the sebaceous follicles of the skin, and is very common in the face, more especially when the seat of acne. In the follicles of the nose they are present in the majority of living persons, and, according to



Simon, are almost universal in dead bodies. He frequently found them living six days after the death of the individual in whom they were found. The animal measures from 1-135th to 1-64th of an inch in length, and from 1-155th to 1-555th of an inch in breadth. It is composed of a head, a thorax, and abdomen.

The head represents in form a truncated cone, flattened


Fig. 97. Cul-de-sac of a sebaceous follicle, containing three animalcules in different positions, and two eggs—(after Gruby).

350 diam.

from above downwards, and directed obliquely downwards

from the anterior part of the trunk. The existence of an eye has not been determined. The head is furnished with two maxiliary palpi, which admit of extensive motion. The thorax is the broadest part of the animal, and is composed of four segments. each of these, on each side, are the legs, eight in number. The abdomen varies in length, is annulated in structure, admitting of certain movements. Internally, Dr. Erasmus Wilson has traced out an alimentary canal, and its termination in an anus, together with a brownish mass he considers to be the liver. No sexual differences have been discovered in them, and they possess no respiratory organs.

The animalcule is easily found by compressing with two fingers the skin we wish to examine, until the sebaceous matter is squeezed out, in the form of a little worm. matter should be placed in a



drop of oil previously heated, then separated with needles,


Fig. 98. Hair and its follicle, in which may be seen the animalcules descending towards the root of the hair, and cul-de-sac of the follicles—(after Gruby).

100 diam.

and examined with a microscope magnifying 250 diameters. Their movements are slow, whilst the conformation of their articulations only permits them to move forwards and backwards, like lobsters. (Gruby.) They are nourished by the sebaceous secretion of the follicles.

They most commonly occupy the excretory duct of the follicles, which are often dilated in the places where they are lodged. Their head is always directed towards the base of the gland. When there are many together, they are placed back to back, and their feet are applied against the walls of the duct. When very numerous, they are compressed closely together, and are found deeper in the ducts. They rarely exist, however, at the base of the gland. In young persons they generally vary in number from two to four; in an aged individual, they may be from ten to twenty. (Gruby.)

2. Cutaneous Ulcers.—In healthy granulating sores, whilst the surface is covered with normal pus corpuscles (Fig. 43), the granulations themselves present fibre-cells in all stages of development passing into fibres. (Fig. 99.) In



scrofulous and unhealthy sores, the pus is more or less broken down, or resembles tubercle corpuscles. (Fig. 46.)

Fig. 99. Fibre cells passing into fibres.

Fig. 100. Fibrous tissue formed from fibre cells.

250 diam.

The epithelial ulcer is very common on the under lip, commencing in the form of a small induration or wart, but rapidly softening in the centre, assumes a cup-shaped depression, with indurated margins, which extend in a circular form more or less over the cheek and chin. An






Fig. 101.

Fig. 102.

examination of the softened matter sometimes exhibits epithelial cells, mingled with fibre or fibro-plastic cells, as in Figure 101. At other times the cells are enlarged, flattened out, and more or less loaded with fat molecules and granules. (Fig. 102.) These growths, though generally denominated cancer, are at once distinguished by a microscopic examination, as may be ascertained by comparing the above figures with groups of cancer-cells. The so-called chimney-sweep's cancer of the scrotum is essentially a similar formation, consisting externally of flattened epithelial scales passing into fibres. (Fig. 103.) Deeper seated, are either groups of younger cells, or concentric layers of aggregated scales, such as those formerly figured in an ulcer of the tongue. (Fig. 28.)

Fig. 101. Epithelial and fibre-cells, from the surface of an ulcer of the lip. Fig. 102. Altered epithelial cells, from the surface of another labial ulcer.

The cancerous ulcer of the skin is often difficult to distinguish microscopically from the epithelial ulcer, because



the external layer, like it, is often composed of softened

Fig. 103. Epidermic scales in mass, and isolated from the chimney-sweep's cancer.

Fig. 104. Group of deep seated-cells.


Fig. 105. The same, after the addition of acetic acid.

250 diam.

epidermis. When, however, a drop of cancerous juice can

be squeezed from the surface, it is found to contain groups of cancer cells, which, from their general appearance, may for the most part be easily distinguished. Considerable experience, however, in the knowledge, and skill in the demonstration, of cancerous and cancroid growths, are necessary in order to pronounce confidently on this point, and to this end an acquaintance with the whole subject of the histology of morbid growths is essential.\*

It may sometimes appear that we can exactly determine the nature of morbid growths which are capable of being reached with the fingers in mucous passages, as in the pharynx, vagina, etc., by the use of the microscope. I have seen several cases where operations have been avoided or undertaken in consequence of the information so arrived at.




On the surface, histological examination is still more easily

\* See the Author's Treatise on Cancerous and Cancroid Growths. Edinburgh, 1849.

Fig. 106. Appearance of section of cancerous ulcer of the skin. a, Epidermic scales and fusiform corpuscles on the external surface. b, Group of epidermic scales. c, Fibro-clastic tissue of the dermis. d, Cancer-cells infiltrated into the fibrous tissue, and filling up the loculi of dermis. 250 diam.

performed, as must be apparent from the position of the soft



cancer of the eye here figured, which was extirpated by Mr. Syme, and recorded by Dr. Dobie.

## USE OF CHEMICAL TESTS.

The chemical examination of urine, blood, milk, and other animal fluids, as well as the detection of poisons in vomited matters, or other organic mixtures and tissues, constitutes an extensive field of inquiry—for a description of which, I must refer to works on chemistry and medical jurisprudence. At the bed-side much of this kind of investigation is now superseded by the use of the microscope,

Fig. 107. a, Encephaloid tumour of the orbit and of the conjunctiva. b, Cancercells in the tumour. c, The same, after the addition of acetic acid.—(after Dobie.)

which at a glance enables us to detect the poverty and adulterations of milk, the spissitude and altered conditions of blood, the nature of various salts and precipitates in urine, etc. The action of chemical re-agents on the corpuscles, made visible by this instrument, has already been alluded to. Chemical tests are most valuable at the bed-side to determine the presence of albumen, bile, sugar, or chlorides in the urine, to which points alone I shall in this place direct your attention.

The Specific Gravity of the Urine is at once obtained by means of a urinometer, and should always be noted at the commencement of the examination of this fluid, as it furnishes important indications for further proceedings. Thus the specific gravity is generally diminished in chronic cases of Bright's disease, and increased in cases of Diabetes.

To detect Albumen in the Urine.—Boil a portion of urine in a test tube over the flame of a spirit lamp, and observe the result. If there be much albumen, the entire fluid becomes white and opaque. If there be a small quantity, a greater or less haziness only may be occasioned. As this latter appearance may be induced by a superabundance of phosphates, nitric acid should be added, when, if the cloudiness be owing to salts, it will disappear, but if to albumen, it will be precipitated. A small quantity of nitric acid added to urine may also occasion haziness, which, however, if owing to the presence of salts, is re-dissolved by an excess of the acid. If both heat and nitric acid cause a white precipitate, albumen is undoubtedly present.

To detect Bile in the Urine.—The test for bile is nitric acid, which changes the fluid containing it in any quantity, first into a grass green, and then, if the test be added in

excess, into a ruby-red or reddish brown tint. If the urine be very much loaded with bile, as sometimes happens in cases of jaundice, so that it resembles porter in appearance, it is better to dilute it with water before adding the acid. If the test be applied to the urine, placed in a clean white plate, so as to form a thin layer over the surface, the play of colours may often be distinctly seen assuming green, violet, pink, and yellow hues. If a very minute quantity of bile only exist in urine, the test produces at once a pink colour.

To detect Sugar in the Uine.—The three best tests for sugar in urine are those known as Moore's test, Trommer's test, and the Fermentation test. Moore's test consists in boiling, for five minutes, urine in a tube, with half its bulk of liquor potassæ. If sugar be present, the liquid assumes a brownish bistre colour. Trommer's test consists in adding one or two drops of a solution of sulphate of copper, so as to give the urine a pale blue colour. Liquor potassæ is then added to the extent of half the volume of urine employed. A pale blue precipitate of hydrated oxide of copper is thrown down, which, if sugar be present, is immediately dissolved, forming a purplish blue solution. The mixture now is to be boiled for a few minutes, when the presence of sugar is indicated by the precipitation of a reddish or yellowish-brown precipitate of suboxide of copper. If no sugar be present, the common oxide of copper is thrown down as a black precipitate. Care must be taken not to add too much sulphate of copper in the first instance. Fermentation test.—A few drops of yeast should be added to urine, and a test tube completely filled with the mixture, should be inverted and allowed to remain in a saucer, containing a little more of the urine. The whole should then be put in a warm place, of about 70 or 80 degrees, for 24 hours. Fermentation

ensues, and carbonic acid is formed, which collects at the top of the tube displaying the fluid. This is a valuable test, because it betrays the existence of a very minute quantity of sugar in the urine, and because, by using a graduated tube of a determinate bulk, the amount of the sugar can be calculated. Two equivalents of carbonic acid are equal to one equivalent of sugar.

The following solution kept prepared is very useful when many observations are made to detect grape sugar in the urine. Take of bitartrate of potash and crystallized carbonate of soda, of each 150 parts, of caustic potash 80 parts, of sulphate of copper 50 parts, and of water 1000 parts; dissolve the carbonate of soda and potash in part of the water boiling, then add the sulphate of copper powdered. When all the bitartrate is dissolved, add the rest of the water, and filter. A few drops of this solution added to a little urine in a test tube will, under the action of heat, throw down a dirty green or yellow precipitate of sub-oxide of copper, if sugar be present.

To detect Chlorides in the Urine.—Add to urine in a test tube an equal volume of distilled water to hold the salts in solution—then add a small drop of nitric acid to neutralize any ammonia that may be present; lastly, add a drop of a solution of nitrate of silver, when a dense white precipitate of the chlorides is precipitated. This is again soluble in an excess of liquor ammoniæ. If there be no chlorides, the urine remains clear, or if they be diminished in quantity, it becomes, on adding the nitrate of silver, more or less hazy.

In concluding this subject allow me to impress upon you the great importance of making yourselves acquainted with all the modes of examination I have described, rather than with one or more of them. It too frequently happens that

exclusive attention to a particular method of exploration, has rendered some medical men good observers of symptoms. whilst they are unacquainted with physical diagnosis; and again, among those who have cultivated the latter, there are some who can percuss and employ the stethoscope with skill, who are ignorant of the use of the microscope, and the contrary. Now you should regard all instruments only as means to an end. In themselves they are nothing, and can no more confer the power of observing, reflecting, or of advancing your knowledge of any disease than a cutting instrument can give the judgment and skill necessary for performing a great operation. We should learn to distinguish between the mechanical means necessary for arriving at truths, and those powers of observation and mental processes which enable us to recognise, compare, and arrange the truths themselves. At the same time, these means are absolutely necessary in order to arrive at the facts on which all correct reasoning is based; and perhaps no kind of knowledge has been so much advanced in modern times by the introduction of instruments and by physical means of investigation, as that of medicine. These enable the practitioner to extend the limits to which otherwise his senses would be limited. Pleximeters, stethoscopes, microscopes, specula. probes, chemical tests, etc. etc., are all useful, and in particular cases indispensable. I do not say employ one to the exclusion of the others, but be equally dexterous in the use of each. Do not endeavour to gain a reputation as a microscopist, as a stethoscopist, or as a chemist; but by the appropriate application of every instrument and method of research, seek to arrive at the most exact diagnosis and knowledge of disease, so as to earn for yourselves the title of enlightened medical practitioners. Above all do not be led away by the notion that any kind of reasoning or theory will enable you to dispense with the careful observation of

facts. What is called tact and skill in detecting maladies is not a peculiar intuition, or a superior power of intelligence possessed by certain persons, but is the result of constant and laborious examination of symptoms and signs in the living, combined with careful research into the nature of morbid organic changes discovered in the dead.

## LECTURE VI.

# THE CLASSIFICATION AND DIAGNOSIS OF CUTANEOUS DISEASES.

Notwithstanding the great advances which have been made in our knowledge of diseases of the skin, it cannot be denied that very inexact notions prevail regarding this class of disorders. I do not here allude to the eruptive fevers, which, from their frequency and danger, necessarily demand the attention of every professional man, so much as to the lighter and more chronic disorders to which the skin is subject. Ignorance, however, here, although it seldom occasions danger to human life produces great inconveniences, exasperates the progress of other maladies, renders life miserable, and frequently destroys those social relations and ties which constitute happiness.

A lady was seized with an eruption on the genital organs, which rendered the slightest contact unbearable. Her husband suspected that she laboured under syphilis, and accused her of infidelity. A medical man, who was consulted, pronounced her disease venereal,—a separation took place between the parties; the lady always maintaining her innocence, but anxious to escape the unfounded suspicions and ill-treatment of her husband. Mercury and an antivenereal treatment were continued for some time, but the disease increased in intensity. At length another physician, skilled in the diagnosis of skin diseases, was consulted, who

pronounced it to be an eczema rubrum, quite unconnected with syphilis; and, on the application of appropriate remedies, a speedy cure confirmed his diagnosis.

A lady in the country sent one of her servants into town, to obtain advice for an eruption which had broken out on her body, and which she was afraid might be communicated to her children. The practitioner consulted was much puzzled, and asked me to see the patient, who, according to him, was labouring under a rare form of skin disease. I found a herpes zoster extending round one half the trunk, and told him it would disappear spontaneously in a few days, which it did.

Nothing is more common in practice than to meet with cases among servants, where prurigo has been mistaken for itch, causing great alarm to the family, and much injury to the servant. The various diseases of the scalp also are continually confounded together. Indeed, examples might easily be accumulated, proving the inconvenience which an unacquaintance with skin diseases may occasion both to patient and practitioner. A young medical man is especially liable to be consulted in cases of trifling skin eruptions; and nothing is likely to establish his credit, as the ready diagnosis and skilful management of such disorders, especially when (as frequently happens) they have been of long standing, and baffled the efforts of older practitioners. Conceiving, then, that this subject deserves more careful consideration than it usually meets with in a clinical course, I propose directing your attention to the classification and general diagnosis of these disorders, as a further introduction to the study of individual cases in the wards.

### CLASSIFICATION.

Skin diseases are so various in appearance and in their nature, that many experienced practitioners have endea-

voured to facilitate their study by arranging them in groups.

There are three kinds of classification which deserve notice:—1st, The artificial classification of Willan; 2d, The natural arrangement of Alibert; and 3d, A pathological arrangement founded on the supposed morbid lesions.

Of these, the best, and the one which most facilitates the study of cutaneous diseases, is certainly that of Willan. No doubt it has its faults and inconveniences, but many of them have been removed by Biett. This classification is founded upon the character presented by the eruption, which, when once known, determines the disease. It is an old saying, that it is much easier to play the critic and to find fault, than to construct something better. This remark may be well applied to those who have ventured to set aside Willan's arrangement, and bring forward others. natural classification of Alibert can never be followed by the student, and presupposes a considerable knowledge of the subject. The pathological arrangement again is decidedly faulty. The morbid anatomy and pathology of many skin diseases are unknown; how, then, can we found a classification upon them? Indeed, the very principles on which such classifications are based, are continually undergoing changes as pathology advances.

On the whole, therefore, the arrangement best suited to the student and for practical purposes is that of Willan and Bateman, with the modifications introduced into it by M. Biett.

Definitions.—Before we can proceed to refer any particular disease to its appropriate class, we must be acquainted with the characteristic appearances which distinguish the different orders. They are as follows:—

1. Exanthema (Rash). — Variously formed irregular sized superficial red patches, which disappear under pressure, and terminate in desquamation.

- 2. Vesicula (Vesicle).—A small, acuminated, or orbicular elevation of the cuticle, containing lymph, which, at first clear and colourless, becomes often opaque or pearl-coloured. It is succeeded either by scurf or a laminated scab.
- 3. Bulla (Bleb).—This differs from the vesicle in its size, a large portion of the cuticle being detached from the skin by the interposition of a watery fluid, usually transparent.
- 4. Pustula (Pustule).—A circumscribed elevation of the cuticle, containing pus. It is succeeded by an elevated scab, which may or may not be followed by a cicatrix.
- 5. Papula (Pimple).—A small, solid, acuminated elevation of the cuticle, in appearance an enlarged papilla of the skin, commonly terminating in scurf, and sometimes, though seldom, in slight ulceration of its summit.
- 6. Squama (Scale).—A lamina of morbid cuticle, hard, thickened, whitish, and opaque, covering either small papular red elevations, or larger deep-red, dry surfaces.
- 7. Tubercula (Tubercle).—A small, hard, indolent, primary elevation of the skin, sometimes suppurating partially, sometimes ulcerating at its summit.
- 8. Macula (Spot).—A permanent discoloration of some portion of the skin, often with a change of its structure. These stains may be white or dark-coloured.

The different appearances thus described characterise the eight orders of Willan and Bateman, viz., 1. Exanthemata; 2. Vesiculæ; 3. Bullæ; 4. Pustulæ; 5. Papulæ; 6. Squamæ; 7. Tuberculæ; 8. Maculæ. The principal modifications made by Biett are removing from these groups certain diseases which have no affinity with them, and forming them into extra orders of themselves. Thus he makes altogether fifteen orders, as seen in the following classification given by his pupils Schedel and Cazenave, which also indicate the subdivisions into which each order is divided:-

ORDER I Exanthemate	z. Vaccinia.	Frambæsia.
Rubeola.	Ecthyma.	Cheloidea.
Scarlatina.	Impetigo.	ORDER VIII.—Maculæ.
Erythema.	Acne.	Lentigo.
Erysipelas.	Mentagra.	Ephelides.
Roseola.	Porrigo.	Nævi and Vitiligo.
Urticaria.	Equinia.	ORDER IX.—Purpura.
ORDER II.—Vesiculæ.	ORDER V.—Papulæ.	X.—Pellagra.
Eczema.	Lichen.	XI.—Radesyge.
Herpes.	Prurigo.	XII.—Lepra Astracha-
Scabies.	ORDER VI.—Squamæ.	nica.
Miliaria.	Psoriasis.	XIII.—The Aleppo
Varicella.	Pityriasis.	Evil, or Malum
ORDER III.—Bullæ.	Ichthyosis.	Alepporum.
Pemphigus.	ORDER VII.—Tubercul	æ. XIV. — Elephantiasis
Rupia.	Lepra Tubercu	losa. Arabica.
ORDER IV.—Pustulæ.	Lupus.	XV. — Syphilidæ or
Variola.	Molluscum.	Syphilitic Eruptions.

Even this classification is very complicated, and appears to me to admit of still further modifications, which will render the subject more simple and practical at the bed-side. I shall point out to you, in the first instance, the reasons which have induced me to make these modifications, and then give, in a tabular form, the classification which we shall in future adopt.

In the orders Exanthemata and Pustulæ, we find several diseases which are characterised by excessive fever, so that they have long been spoken of under the term of eruptive fevers, as well as under that of febrile eruptions. With them, in short, fever is the characteristic, and they are influenced by laws of a peculiar character, altogether different from those which regulate the production of other cutaneous affections. I propose, then, removing these disorders from the category of skin diseases altogether, which will only leave three in the first order, namely, erythema, roseola, and urticaria. I am aware that, strictly speaking, these may be accompanied by slight fever, which may also occur in several other skin diseases. But I do not pretend to form a classification which is perfect, or even pathological, but one which some experience in the teaching

of these diseases has convinced me is useful and practical for the student.

In the order *Vesiculæ* we find five diseases. I propose cutting out miliaria, as being very unimportant, and a trifling sequela of fevers. Varicella I believe to be a modified small-pox, and I omit it for the same reasons as I do variola. Scabies, on the other hand, though dependent upon the presence of an insect, the *Acarus Scabiei*, presents such distinct characters as to warrant its retention.

I propose expunging the order Bullæ altogether. We find in it two diseases. The first of these, pemphigus or pompholyx, is a vesicular disease in every point, appearing in successive crops, and forming a laminated scab. Rupia, on the other hand, is evidently a pustular disease, forming a prominent scab, producing ulceration, and leaving a cicatrix. I shall, therefore, add pemphigus to the order vesiculæ, and rupia to that of the pustulæ.

From the Pustulæ, for the reasons formerly stated, I expunge variola, vaccinia, and equinia. Mentagra, so far as I have been able to study it in this country, has always consisted of eczema or impetigo on the chin of the male. In syphilitic cases, it is more or less tubercular, and it has been described also as consisting of a vegetable parasite. Although I have never seen the appearance figured by Cazenave (Plate 16), I can understand that such a mentagra might really consist of vegetable fungi. At all events mentagra is not a special pustular disease. Porrigo means any eruption on the head, whether vesicular, pustular, or squamous. Favus, to which it has long been applied, is undoubtedly a vegetable parasite, and ought, with others of a like nature, to constitute a class of themselves. Moreover, it is neither vesicular nor pustular. Hence the class of pustulæ will with us contain only impetigo, ecthyma, acne, and rupia.

The orders Papulæ and Squamæ remain the same. The strophulus of many English writers is certainly only lichen

occurring in the child; and what has been called lepra, as distinguished from psoriasis, is the latter disease presenting an annular form.

From the class *Tuberculæ* I cut out frambæsia, as being a disease unknown in this country, together with cheloidea, which, as I understand it, means either cancer or tubercle of the skin.

As regards the order *Maculæ*, I place in it, as did Willan, purpura, because, although sometimes it may depend on constitutional causes of an obscure nature, and at others be allied to scurvy, it still, in an arbitrary classification of this kind, constitutes an undoubted spot or macula.

All the other orders of Biett I shall take the liberty of expunging—pellagra, lepra Astrachanica, and malum Alepporum, are unknown in this country. I agree with Hebra in thinking that radesyge is only a modified form of lupus. The elephantiasis Arabica is an hypertrophy of the areolar tissue or chorion, and belongs more to the subject of fibrous growths than that of skin diseases. Syphilitic diseases I do not regard as a distinct order, but as any of the ordinary skin affections, more or less modified by a peculiar state of the constitution.

Whilst I have cut out many diseases from the eight orders originally established by Willan, and subsequently modified by Biett, I find it necessary to add two orders, which the advance of pathology and histology shows ought to be considered apart. I allude to those which depend on the presence of parasitic animals and plants, and which may be called respectively Dermatozoa and Dermatophyta. It has now been shown by M. Bourguignon, that scabies is dependent on the presence of an acarus, but that the insect is only indirectly the cause of the eruption. Hence I put acarus among the dermatozoa, although it certainly forms, when present, a constituent of itch. Among the dermatophytes will be placed favus and mentagra,—both

removed from the class pustulæ. Other diseases, such as plica Polonica, and pityriasis, have been considered as parasitic, but the former is unknown in this country, and the latter, though frequently presenting epiphytes among the scales, owes none of its essential characters to this circumstance.

The classification, then, we shall in future adopt is as follows:—

ORDER I.—Exanthemata.	ORDER IV.—Papulæ.	Nævi.
Erythema.	Lichen.	Purpura.
Roseola.	Prurigo.	ORDER VIII.—Dermatozoa.
Urticaria.	ORDER VSquamæ.	Entozoon folliculo-
ORDER II Vesiculæ.	Psoriasis.	rum.
Eczema.	Pityriasis.	Acarus.
Herpes.	Ichthyosis.	Pediculus.
Scabies.	ORDER VI.—Tuberculæ. ORDER IX.—Dermatophy-	
Pemphigus.	Lepra Tuberculo	sa, tæ.
ORDER III.—Pustulæ.	Lupus.	Achorion Schönleini
Impetigo.	Molluscum.	(Favus).
Ecthyma.	ORDER VII.—Maculæ.	Achorion Grubii
Acne.	Lentigo.	(Mentagra).*
Rupia.	Ephelides.	

#### DIAGNOSIS.

The recognition of skin diseases, and the separating one class from the other, is of essential importance to a proper treatment. On this point I fully agree with a recent writer, who says "the treatment of a great many cutaneous diseases is but of secondary importance, compared with their differential diagnosis. Many of them will get well without

<sup>\*</sup> It has been objected to the words porrigophyte and mentagraphyte, introduced by Gruby, that they are unclassical; and, as the celebrated botanist Link, after carefully examining these vegetations, has described the former as a new genus, under the head of achorion (from achor, the old term given to a favus crust by Willan), I have thought it best to adopt that term. To mark the variety in favus, he has added the name of its discoverer, Schönlein; and I have ventured, at all events provisonally, to distinguish the one described as existing in mentagra, by adding to it, also that of its discoverer, Gruby.

any treatment, provided they are allowed to pursue their natural course; and, on the contrary, a mild and simple eruption by being mistaken, from a similarity of external appearances, for one of a severe or rebellious character, and treated accordingly, may be aggravated and prolonged for an indefinite period." (Burgess.) This differential diagnosis, however, to the inexperienced, is a matter of great difficulty, because considerable tact is often necessary, not only to discover the original element each disease presents, such as a rash, vesicle, pustule, scale, and so on, but this is often impossible. Under such circumstances the diagnosis is frequently derived from the scab, or other appearances presented, such as the cicatrix. The whole subject has been rendered very confused and complicated by systematic writers, who have often given different names to the same disease, or unnecessarily divided them into forms and varieties. I advise you not to pay any attention to these forms and varieties for the present, and confine your efforts only to the detection of the diseases enumerated in the table under each order; and with a view of facilitating your endeavours, the following short diagnostic characters and definitions should be attended to.

### I. EXANTHEMATA.

- 1. Erythema.—A slight continuous redness of the skin in patches of various shapes and sizes.
- 2. Roseola.—Circumscribed rose-red patches, of a circular, serrated, or annular form.
- 3. Urticaria.—Prominent red patches of irregular form, the centre of which is often paler than the surrounding skin.

## II. VESICULÆ.

Eczema.—Very minute vesicles in patches, presenting a shining appearance, yielding a fluid which dries into a lami-

nated or furfuraceous crust. The skin is of a bright red colour.

Herpes.—Clusters of vesicles, varying in size from a millet seed to that of a pea, surrounded by a bright red areola. They yield a fluid which dries into a thin incrustation, that drops off between the eighth and fifteenth day.

Scabies.—Isolated vesicles of an acuminated form, commonly seated between the fingers, and flexor surfaces of the arms and abdomen—never on the face.

Pemphigus.—Large vesicles or blebs (bullæ), surrounded by an erythematous circle, the fluid of which forms, when dry, a laminated crust. When chronic, they appear in successive crops, and the disease is called pompholyx.

#### III. PUSTULÆ.

Impetigo. — Small pustules, commonly occurring in groups, and forming an elevated crust.

Ecthyma.—Large isolated pustules, depressed or umbilicated in the centre, and leaving a cicatrix.

Acne.—Isolated pustules situated on a hardened base, which form and disappear slowly. They only occur on the face and shoulders.

Rupia.—Large pustules, followed by thick prominent crusts, and producing ulcerations of various depths.

## IV. PAPULÆ.

Lichen.—Minute papulæ occurring in clusters or patches.

Prurigo.—Larger and isolated papulæ generally seated on the extensor surfaces of the body.

## V. SQUAMÆ. .

Psoriasis.—Whitish laminated scales slightly raised above the reddened surface of the skin. Lepra is psoriasis occurring in rings.

Pityriasis.—Very minute scales, like those of bran, seated on a reddened surface.

Ichthyosis.—Induration of the epidermis, and formation of square or angular prominences, not seated on a reddened surface.

#### VI. TUBERCULÆ.

Lepra Tuberculosa.—(Elephantiasis of the Greeks.)— Tubercles varying in size, preceded by erythema and increased sensibility of the skin, and followed by ulceration of their summits.

Lupus.—Induration or tubercular swelling of the skin, which may or may not ulcerate. In the former case, ulceration may occur at the summit or at the base of the tubercles, and frequently extends in the form of a circle more or less complete.

Molluscum.—Pedunculated, globular, or flattish tubercles, accompanied by no erythema or increased sensibility, occurring in groups. They are filled with atheromatous matter.

## VII, MACULÆ.

Lentigo or Freckle.—Brownish-yellow or fawn-coloured spots on the face, bosom, hands, or neck.

Ephelis.—Large patches of a yellowish-brown colour, accompanied by slight desquamation of the cuticle.

Nævi or Moles.—Spots of various colours and forms, sometimes elevated above the skin. They are congenital.

Purpura.—Red or claret-coloured spots or patches, which do not disappear under pressure of the finger.

### VIII. DERMATOZOA.

These minute animals require a lens of considerable power to ascertain their characters, which need not be particularised here, as they have been previously described and figured in the last lecture. (See p. 114, et seq.)

#### IX. DERMATOPHYTE.

These minute plants also require a high magnifying power to distinguish them. But they communicate peculiar characters to certain cutaneous diseases, as follows:—

Favus.—Bright yellow, umbilicated crusts, surrounding individual hairs, which agglomerate together to form an elevated friable crust, of a peculiar musty or mousey smell. (See pp. 112, 113.)

Mentagra.—Grayish or yellowish dry crusts, of irregular form, originating in the hair follicles of the beard.

In forming your diagnosis, therefore, you will be guided principally by three characters:—1st, The primitive and essential appearance—that is, whether a rash, vesicle, pustule, and so on. 2d, The crust,—whether laminated or prominent, composed of epidermis only, etc. 3d, Ulceration—whether present or absent; and if so, the kind of cicatrix. These and other characters I shall point out at the bedside, so as to familiarise you with their appearances.

You will remember that the classification formed by Willan is wholly artificial. It is like the Linnæan classification of plants. The difficulty for the learner is to recognise the essential character, the more so as many diseases pass through various stages before this is formed. Thus herpes presents, 1st, a rash; 2d, papules; 3d, vesicles; 4th, pustules; yet the disease is considered vesicular. Ecthyma passes through the same stages, yet it is considered pustular. In the vesicular disease, however, the crust is laminated,—in the pustular, it is more or less prominent.

Again, it not unfrequently happens that two or more diseases are combined together in one eruption. Thus it is

very common to meet eczema and impetigo combined, when the disease is called *Eczema-Impetiginodes*. Favus occasionally causes considerable irritation, producing a pustular or impetiginous margin around it. The vesicles of scabies are often accompanied by the pustules of ecthyma, and so on.

In very chronic skin diseases, it may happen that it is impossible to say what the original disorder was, whether vesicular, pustular, scaly, or papular. In such cases the skin assumes a red colour, the dermis is thickened, the epidermis rough and indurated, and a morbid state is occasioned, in which all trace of the original disease is lost, and what remains is a condition common to various disorders.

As regards varieties, little need be said, and as formerly stated, I advise you to postpone their study until you are acquainted with the diseases themselves. Even then an acquaintance with them is of secondary importance. These varieties have been formed on account of the most varied circumstances, such as,—1st, Duration, most of them may be acute or chronic; 2d, Obstinacy, hence the terms fugax, inveterata, agrius, etc.; 3d, Intensity, hence the terms mitis, maligna, etc.; 4th, SITUATION, hence the terms capitis, facialis, labialis, palmaris, etc.; 5th, Form, hence the terms circinatus, scutulata, iris, gyrata, larvalis, figurata, tuberosa, quttata, etc.; 6th, Constitution, hence the terms cachectica, scorbutica, syphilitica, etc.; 7th, Age, hence the terms infantilis, senilis, etc.; 8th, Colour, hence the terms album, nigrum, rubrum, versicolor, etc.; 9th, Density, hence the terms sparsa, diffusa, concentricus, etc.; 10th, Feel, hence the terms læve, indurata; 11th, Sensation Produced, hence the terms formicans, pruritus, urticans, etc.; 12th, Geogra-PHICAL DISTRIBUTION, hence the terms tropicus, Ægyptiana, Norwegiana, etc.

#### Porrigo.

There was a period in the history of skin diseases when they were arranged in two great divisions, viz., those affecting the scalp, and those affecting the rest of the cutaneous surface. All the disorders comprehended in the first of these divisions received the name of Porrigo, a word, said by some to be derived from porrum, on account of the scales or concretions of the scalp, resembling the layers of an onion; by others, it is derived from porrigo, to spread. Willan described six kinds of Porrigo, viz., P. larvalis. P. furfurans, P. scutulata, P. favosa, P. lupinosa, and P. decalvans. It is now ascertained that none of these diseases are necessarily peculiar to the scalp,—and that, although they are more or less modified by being connected with and affecting the hairs of that region, they may also occur on other parts of the skin. There can be little doubt, however, that the employment of the term Porrigo, as well as the corresponding word Teigne in France, has thrown great confusion over the subject of eruptions on the scalp. But, as this term is still in pretty general use, it will be well to explain to you what diseases these different kinds of Porrigo really are.

Porrigo larvalis (larva, a mask) is really Impetige, or Eczema impetiginodes, of the scalp. The former is recognised by crusts more or less prominent or nodulated; the latter, by the circumstance that, in addition to these nodules, there is between them a laminated or brittle crust, spread more or less equally over the surface. They are both very common in infants and children; and, as the disease sometimes extends over the face, concealing the features, hence the term larvalis. A very characteristic representation of impetigo capitis, which I now show you, is given in Willan and Bateman, Plate xli., erroneously called Porrigo Favosa.

See also the disease on the face. *Ibid.*, Plate xxxvii. *Alibert* Planches 13 and 15.

Porrigo furfurans (furfur, bran) is really Pityriasis of the scalp, although Psoriasis of that region has also received the same appellation. There is also a peculiar form of Eczema, or Eczema impetiginodes, in which the crust is friable, and breaks up, or crumbles into minute fragments, to which the term furfurans has been erroneously applied. The true Porrigo furfurans (Pityriasis) is well represented. — Willan and Bateman, Plate xxxviii. Alibert, Planches 14 and 15.

Porrigo scutulata (scutulum, a small shield).—The nature of this disease has been much disputed. By some, it is said to be Favus (Erasmus Wilson), by others a form of Herpes (Cazenave). The disease is described by Willan and Bateman, and more recently by Burgess, to consist of oval or rounded, slightly elevated patches, covered with furfur, and having stunted or filamentous hair projecting from the surface. It is a form of skin eruption exceedingly rare in Edinburgh. It seems to be represented, Willan and Bateman, Plate xxxix. Willis (Trichosis Scutulata).

Porrigo favosa (favus, a honeycomb) is a disease, the true nature of which has been only lately determined. It consists essentially in an exudation on the skin, in which fungi or phytaceous plants grow. Round, isolated, bright yellow crusts are formed, which, when compressed together, assume an hexagonal shape. Hence the term favosa. It is well represented, Willis (Trichosis lupinosa). Erasmus Wilson, Fasiculus I. Alibert, Planche 17.

Porrigo lupinosa (lupinum, the lupine).—This is the same disease as the last. The round or oval crusts when

isolated, and at an early stage, present a concavity and form, resembling that of the lupine seed—hence its name.

Porrigo decalvans (calvus, bald).—Baldness is so common among the aged, that it can scarcely be called a disease; but when it occurs in young persons, and is circumscribed, it constitutes the Porrigo decalvans of Willan. It is said by Gruby to depend on a vegetable parasite growing in the hair. It is well represented, Willan and Bateman, Plate xl. Willis (Trichosis Decalvans).

From this analysis of the different kinds of the so-called Porrigo, you observe that there is nothing peculiar with regard to them. With the exception of baldness, none essentially belong to the hairy scalp. True favus is far more common on the head than elsewhere; but I have frequently seen it on various parts of the cutaneous surface, and occasionally on the cheeks or shoulders, without being on the scalp at all. It follows that instead of the term Porrigo, you should designate the disease as Eczema Impetigo, Pityriasis, Psoriasis, or Favus of the scalp, as the case may be.

Notwithstanding I have endeavoured to place this subject before you in as simple and uncomplicated a form as possible, I am conscious that at first you will still experience considerable difficulty in the diagnosis of skin affections. This can only be removed by practical experience at the bed-side, and by constantly exercising your powers of observation in detecting the essential elements which their varied forms present. At the same time, I think the modified classification and short characters I have given, will materially assist your studies in this important department of practical medicine. It must be remembered, however, that they only refer to those cutaneous diseases which you are

liable to meet with in this country. Should you ever be called upon to practise in the tropics, or in other places where peculiar skin disorders prevail, it will, of course, be your duty to study them in an especial manner. Here, as they cannot be made the subject of clinical observation, they are altogether removed from our consideration.

THE END.

#### BY THE SAME AUTHOR.

In Demy 8vo, price 12s.,

## ON CANCEROUS AND CANCROID GROWTHS.

#### CONTENTS.

PART I.—Fifty-six Cases of Can- | Chap. III. General Anatomy. cerous and Cancroid Growths, with Microscopic Examinations, and exact Drawings of the Morbid Structures in each. PART II.—Chap. I. Histology.

II. Chemistry.

IV. General Pathology.

V. Statistics.

VI. Diagnosis. VII. Prognosis.

VIII. Rational Treatment.

With ONE HUNDRED and NINETY ILLUSTRATIONS, copied from Nature, and Drawn on Wood, by the Author.

- "We have seldom, in the course of our reading, met with a book possessing more real interest, novelty, and instruction. We would compare this work, in reference to the use of the microscope in disease, with those of Lænnec or Latham in regard to the use of the stethoscope in thoracic disease."—Medical Gazette.
- "It is a serious and laborious investigation, in which exact research, acute criticism, and great power of generalisation will be found united."—Lancet.
- "The most useful and complete work on cancer which has appeared in our language."—Medical Times.
- "Dr. Bennett's work, we have no doubt, will constitute the standard treatise on the subject; while, to every one engaged in microscopic researches, the volume is indispensable."—Dublin Medical Press.
- "The chapter on treatment, though short, contains many new and important observations."—Dublin Quarterly Journal.
- "Dr. Bennett has added much to our knowledge of cancer, and done still more to clear the path for the observers who will succeed him."—Brit. and For. Med.-Chir. Review.
- " Dr. Bennett's numerous and interesting cases, his careful microscopic examinations, and his graphic illustrations, should not merely be perused, but carefully studied, by every physician and surgeon in connection with his lucid descriptions of each morbid growth, as exhibited in the pages of his very original and able work."—Copland's Medical Dictionary.

Edinburgh: SUTHERLAND & KNOX. London: SIMPKIN, MARSHALL, & Co.

#### BY THE SAME AUTHOR.

Demy 8vo, price 6s., Woodcuts,

- THE PATHOLOGY AND TREATMENT OF PULMONARY TUBERCULOSIS; and on the Local Medication of Pharyngeal and Laryngeal Diseases frequently mistaken for, or associated with, Phthisis. By John Hughes Bennett, M.D., F.R.S.E., Professor of the Institutes of Medicine and of Clinical Medicine in the University of Edinburgh, etc.
- "I have no hesitation in declaring my opinion, that Dr. Hughes Bennett has advanced the farthest in this investigation."—Ancell on Tuberculosis.
- "The Author's views respecting the treatment of Phthisis appear to us to be legitimate deductions from the large body of facts which he has collected and analysed."—Dublin Medical Press.
- "Dr. Bennett is one of the advanced party in medicine, and a deep and lasting debt of gratitude mankind owes him for this volume."—

  Medical Circular.
- "To those who still follow the old palliative system of treating Phthisis, to all who wish to understand the most modern doctrines of the pathology of the disease, and to all who are desirous of studying the principles upon which cod-liver oil is exhibited in its treatment, we can recommend a careful perusal of Dr. Bennett's work."—Medical Times and Gazette.
- "The cases recorded in this volume will prove deeply interesting to the pathologist, many of them being detailed instances of arrested phthisis, upon which the author's views of the curability of the disease have been founded."—Association Medical Journal.
- "Dr. Bennett's late treatise on "Tuberculosis" will of course be read by everybody.—No one can read the work without profit."—British and Foreign Medical Chirurgical Review.
- "To say that we have been pleased with the work would be too faint praise; we have been more than pleased; it has deeply interested us, and has elucidated some points hitherto obscure, and given us a pathology of tuberculosis, on the whole more satisfactory than any we have met with."—Philadelphia Medical Examiner.
- "To Dr. J. H. Bennett belongs the high honour of having clearly pointed out the remarkable efficacy of cod-liver oil, in tubercular diseases."—Headland on the Action of Medicines.
- "If the author had only reported these cases, he would have benefited his profession, and deserved our thanks. As it is, however, his whole volume is so replete with valuable matter, that we feel bound to recommend our readers, one and all, to peruse it."—Lancet.

Edinburgh: Sutherland & Knox. London: Simpkin, Marshall, & Co.



# ENCYCLOPÆDIA BRITANNICA,

## EIGHTH EDITION.

Now Publishing in Monthly Parts, price 8s, and Quarterly Volumes, price 24s.

#### LIST OF SOME OF THE CONTRIBUTORS.

Rt. Hon. Thomas Babington Macaulay.

Rt. Rev. RICHARD WHATELY, D.D., Archbishop of Dublin.

Rt. Rev. R. Dickson Hampden, D.D., Bishop of Hereford.

WILLIAM WHEWELL, D.D., Trinity College, Cambridge.

Sir David Brewster, K.H., LL.D., Principal of the United Colleges of St Salvator and St Leonard, St Andrews.

RICHARD OWEN, Esq., F.R.S.

JOHN LEE, D.D., Principal of the University of Edinburgh.

Sir WILLIAM HAMILTON, Bart.

Sir Archibald Alison, Bart.

Sir John Richardson.

Sir JOHN M'NEILL.

HENRY ROGERS, Esq., Author of the "Eclipse of Faith," &c.

ISAAC TAYLOR, Esq., Author of the "Natural History of Enthusiasm," &c.

Rev. Charles Kingsley, Author of "Hypatia," "Westward Ho," &c.

J. D. Forbes, Professor of Natural Philosophy in the University of Edinburgh, &c. &c.

ROBERT STEPHENSON, Esq., M.P., President of the Institution of Civil Engineers, &c.

RICHARD MONCKTON MILNES, Esq., M.P.

HEPWORTH DIXON, Esq.

THEODORE MARTIN, Esq.

COLONEL PORTLOCK, R.M.A., Woolwich.

DAVID MASSON, M.A., Professor of English Literature in University College, London.

PETER CUNNINGHAM, Esq.

J. Y. Simpson, M.D., Professor of Midwifery in the University of Edinburgh.

THOMAS DE QUINCEY, Esq.

R. G. LATHAM, Esq., M.A., M.D., &c.

W. E. AYTOUN, Professor of Rhetoric and Belles Lettres in the University of Edinburgh.

J. R. M'Culloch, Esq., Member of the Institute of France, &c.

AUGUSTUS PETERMANN, Esq., F.R.G.S., &c.

WILLIAM SPALDING, M.A., Professor of Logic and Metaphysics in the University of St. Andrews.

Rev. G. R. GLEIG, M.A, &c.

Rev. J. E. RYLAND, M.A., &c.

Rev. P. Kelland, M.A., Professor of Mathematics in the University of Edinburgh.

Antonio Panizzi, Esq.

THOMAS ANDERSON, M.D., Professor of Chemistry in the University of Glasgow.

Dr. Doran, Author of "Habits and Men," &c.

WILLIAM FAIRBAIRN, Esq., C.E., Manchester.

H. L. Mansel, Reader in Moral and Metaphysical Philosophy, Magdalen College, Oxford.

James Caird, Esq., Author of "English Agriculture in 1850-51," &c.

THOMAS LAYCOCK, M.D., Professor of the Practice of Physic in the University of Edinburgh.

Daniel Wilson, LL.D., F.R.S.A., &c.

LAWRENCE OLIPHANT, Esq., Author of the "Russian Shores of the Black Sea."

GEORGE FARQUHAR GRAHAM, Esq.

JOHN FLEMING, D.D., Professor of Natural Science, New College, Edinburgh.

E. B. Denison, Esq., M.A., Q.C., &c.

Rev. John Cairns, M.A.

J. H. BALFOUR, M.D., Professor of Botany in the University of Edinburgh.

J. S. Blackie, Professor of Greek in the University of Edinburgh.

JAMES MONTGOMERY, Esq., Author of "Greenland and other Poems," &c.
WILLIAM GREGORY, M.D., Professor of Chemistry in the

University of Edinburgh.

GEORGE WILSON, M.D., Professor of Technology in the University of Edinburgh.

Rev. WILLIAM SCORESBY, LL.D., &c.

ROBERT CARRUTHERS, Esq.

JOHN HILL BURTON, Esq.

J. B. Jukes, Esq., M.A., Vice-President of the Geological Society of Dublin.

WILLIAM HOSKING, Professor of Arts and Construction, King's College, London.

ALEXANDER SMITH, Esq., Author of "A Life Drama," &c.

Rev. WILLIAM HANNA, LL.D.

Rev. Robert Main, M.A., F.R.A.S., &c.

JOHN TULLOCH, D.D., Primarius Professor of Divinity, St. Andrews.

CHARLES MACLAREN, Esq., F.R.S.E., &c.

J. L. RICARDO, Esq., M.P.

LEONHARD SCHMITZ, LL.D., F.R.S.E., &c.

EDWARD THORNTON, Esq., Author of "Gazetteer of India," &c. George Ferguson, LL.D., Professor of Humanity, King's College, Aberdeen.

JAMES PILLANS, M.A., Professor of Humanity in the University of Edinburgh.

F. CRACE CALVERT, Professor of Chemistry, Royal Institution, Manchester.

ROBERT CHRISTISON, M.D., Professor of Materia Medica in the University of Edinburgh.

R. S. Poole, Esq., M.R.S.L., &c.

JOHN BARROW, Esq., Admiralty.

JOSEPH D. HOOKER, M.D., R.N.

CHARLES TOMLINSON, Esq., Editor of a "Cyclopædia of Useful Arts," &c.

W. L. ALEXANDER, D.D.

THOMAS BAZLEY, Esq., Chairman of the Chamber of Commerce, Manchester.

W. H. LANGLEY, Esq., Editor of "Bell's Life in London."

CHARLES MACINTOSH, Esq., Author of the "Book of the Garden."

T. C. ARCHER, Esq., Author of "Popular Economic Botany."

J. P. LACAITA, Esq.

ARTHUR ASHPITEL, Esq.

A. V. KIRWAN, Esq.

WILLIAM BLAIR, Esq.

# ENCYCLOPÆDIA BRITANNICA,

## EIGHTH EDITION

## PRINCIPAL SUBJECTS.

#### VOLUME I.

- DISSERTATION FIRST.—On the Progress of Metaphysical and Ethical Philosophy, since the Revival of Letters in Europe. By Dugald Stewart, late Professor of Moral Philosophy in the University of Edinburgh.
- DISSERTATION SECOND.—On the Progress of Ethical Philosophy, chiefly during the Seventeenth and Eighteenth Centuries. By Sir James Mackintosh, LL.D. With a Preface by William Whewell, D.D.
- DISSERTATION THIRD. On the Rise, Progress, and Corruptions of Christianity. By RICHARD WHATELY, D.D., Archbishop of Dublin.
- DISSERTATION FOURTH.—On the Progress of Mathematical and Physical Science since the Revival of Letters in Europe. By Sir John Playfair, late Professor of Natural Philosophy in the University of Edinburgh.
- DISSERTATION FIFTH.—On the Progress of Mathematical and Physical Science, chiefly during the Eighteenth Century. By Sir John Leslie, late Professor of Natural Philosophy in the University of Edinburgh.
- DISSERTATION SIXTH.—On the Progress of Mathematical and Physical Science during the Nineteenth Century. By JAMES D. FORBES, Professor of Natural Philosophy in the University of Edinburgh.

#### VOLUME II.

- ABERRATION and ACOUSTICS. By George Buchanan, F.R.S.E.
- ACHROMATIC GLASSES and AERONAUTICS. By Sir John Leslie.
- **ADMIRAL and ADMIRALTY.** By Sir John Barrow, Bart Revised by John Barrow, Admiralty.
- **ADDISON**, (JOSEPH.) By WILLIAM SPALDING, A.M., Professor of Logic in the University of St Andrews.
- **ÆSCHYLUS.** By J. S. BLACKIE, Professor of Greek in the University of Edinburgh.
- **ÆTNA** and ALPS. By ROBERT JAMESON, late Professor of Natural History in the University of Edinburgh.
- AFGHANISTAN. By David Buchanan and Edward Thornton, East India House.
- AFRICA. By Augustus Petermann, F.R.G.S.
- AGRICULTURE. By John Wilson, Eddington Mains, Berwickshire.
- AGRICULTURAL CHEMISTRY. By THOMAS ANDERSON, M.D., Professor of Chemistry in the University of Glasgow.
- AGRARIAN LAWS. By George Ferguson, Professor of Humanity, King's College, Aberdeen.
- ALGEBRA. By WILLIAM WALLACE, LL.D., late Professor of Mathematics in the University of Edinburgh.
- AMERICA. By Charles Maclaren, F.R.S.E.
- ANATOMY. By David Craigie, M.D., F.R.S.E.
  - Abbreviation—Abyssinia—Academy—Algiers—Alphabet
    —Alum.

#### VOLUME III.

ANATOMY. By DAVID CRAIGIE, M.D., F.R.S.E.

ANCHOR. By Jonathan Aylen, Greenwich.

ANDES, ANGLE, &c. By Sir John Leslie.

ANGLING, ANIMAL KINGDOM, and ANIMALCULE. By JAMES WILSON, Author of various works on Natural History.

ANNUITIES. By Joshua Milne.

ANT. By P. M. ROGET, M.D., F.R.S.

ANTRIM and ARMAGH. By HENRY SENIOR.

APPARITIONS. By James Browne, LL.D.

AQUEDUCT. By George Buchanan, C.E.

ARABIA. By D. Buchanan and W. Plate, LL.D.

ARCH. By John Robison, late Professor of Natural Philosophy in the University of Edinburgh, and William Wallace, LL.D.

ARCHÆOLOGY. By DANIEL WILSON, LL.D.

**ARCHITECTURE.** By WILLIAM HOSKING, Professor of Arts and Construction, King's College.

ARISTOTLE and ARISTOTLE'S PHILOSOPHY. By R. Dickson Hampden, D.D., Bishop of Hereford.

ARMY. By J. Browne, LL.D., and J. H. STOCQUELER.

ARTILLERY. By Capt. Spearman and Colonel Portlock.

ARTS. By W. HAZLITT and W. B. JOHNSTONE, R.S.A. .

ASIA. By H. MURRAY, F.R.S.E., and W. PLATE, LL.D.

ASSAYING. By ROBERT MUSHET, of the Royal Mint.

ASTRONOMY. By Thos. GALLOWAY, Sir John Playfair, and Thos. Henderson. With Supplement by Rev. Robert Main, Royal Observatory, Greenwich.

Archery-Arithmetic.

### VOLUME IV.

- ASTRONOMY. By Thos. Galloway, Sir John Playfair, and Thos. Henderson. With Supplements by Rev. R. Main, Royal Observatory, Greenwich.
- ATHENS and ATTICA. By JAMES BROWNE, LL.D. Revised by Dr L. Schmitz, F.R.S.E., Rector of the High School of Edinburgh.
- ATMOMETER, BAROMETER, and BAROMETRICAL MEASUREMENTS. By Sir John Leslie. With Supplements.
- ATMOSPHERE. By Thomas Thomson, late Professor of Chemistry in the University of Glasgow.
- ATTERBURY. By the Right Hon. THOMAS BABINGTON MACAULAY.
- ATTRACTION. By James Ivory, F.R.S.
- AURORA BOREALIS. By ROBT. JAMESON, F.R.S., late Professor of Natural History in the University of Edinburgh.
- AUSTRALASIA and AUSTRALIA. By Sir John Barrow. With continuation by Samuel Mossman, Author of the "Gold Fields of Australia," &c.
- AUSTRIA. By EMERIC SZABAD, Author of "Hungary, Past and Present."
- AVERAGE. By JOHN WARRACK, Average Stater.
- BACON. By WILLIAM SPALDING, Professor of Logic in the University of St Andrews.
- BAKING, BLEACHING, &c. By James Stark, M.D., F.R.S.E.
- BAILLIE, BARBOUR, BALLAD, BARCLAY, &c. By DAVID IRVING, LL.D.
- BALANCE OF POWER and BIBLIOGRAPHY. By MACVEY NAPIER, late Professor of Conveyancing in the University of Edinburgh.
- BALLAT, BANKRUPTCY, and BENTHAM. By J. H. Burton, Author of the "History of Scotland," &c.
- BASKET-MAKING. By Sir J. G. DALZELL.
- BATHING, BECCARIA, &c. By Thomas Young, M.D.
- BENGAL. By EDWARD THORNTON, East India House.
- BEAUTY. By LORD JEFFREY.
- BEE. By JAMES WILSON, F.R.S.E.

### VOLUME IV.—Continued.

BEETHOVEN. By George Farquhar Graham.

BELGIUM.

BELL, (SIR CHARLES.) By Sir John M'NEILL.

BIBLE and BIBLE SOCIETIES. By Rev. JAMES TAYLOR, D.D.

BLACK SEA. By LAWRENCE OLIPHANT, Author of "The Russian Shores of the Black Sea."

BLASTING. By ROBERT and THOMAS STEVENSON, Civil Engineers.

BOHEMIAN BRETHREN. By James Montgomery, Author of "Greenland and other Poems."

BOLIVIA.

Babylon—Baptism—Bernouilli—Blind—Block-Machinery—Blowpipe.

#### VOLUME V.

BOOKBINDING. By CHARLES MARTEL.

BOMBAY, BURMAH, &c. By EDWARD THORNTON, East India House.

**BOOK-KEEPING.** By Joseph Lowe.

**BORNEO** and **BORNOU**. Revised by Augustus Petermann, F.R.G.S., &c.

BOTANY. By JOHN HUTTON BALFOUR, M.D., Professor of Botany in the University of Edinburgh.

**BRAHMINS.** By James Browne, LL.D.

BRASS. By Charles Sylvester, C.E.

BRAZIL.

BREAKWATER. By Sir John Barrow, Bart. Revised by John Barrow, Admiralty.

BREWING. By James Stark, M.D., F.R.S.E.

BRICKMAKING. By SAMUEL HOLMES, Liverpool.

BRIDGE. By Thomas Young, M.D., F.R.S.

BRITAIN. By James Browne, LL.D., with continuation.

BRUCKER. By Sir WILLIAM HAMILTON, Bart.

BUILDING. By WILLIAM HOSKING, Author of "Architecture."

BUNYAN. By the Right Hon. THOMAS BABINGTON MACAULAY.

BURMAH. By D. BUCHANAN and E. THORNTON.

## VOLUME VI.

BISHOP BUTLER. By HENRY ROGERS, Author of the "Eclipse of Faith," &c.

BURNING-GLASSES. By George Buchanan, F.R.S.E.

CALENDAR. By Thomas Galloway, F.R.S.

CALIFORNIA.

CALVIN and CHANNING. By Rev. W. L. ALEXANDER, D.D.

CAMPBELL, THOMAS. By W. E. AYTOUN, Professor of Rhetoric and Belles Lettres in the University of Edinburgh.

CANADA. By J. B. Brown, Author of "Views of Canada."

CANARY ISLANDS. By J. Y. Johnson, Madeira.

CANNON. By Colonel PORTLOCK, Woolwich.

CAPILLARY ACTION. By JAMES IVORY, F.R.S.

CARTHAGE. By JAMES BROWNE, LL.D., and Dr SCHMITZ.

CARPENTRY and CHROMATICS. By THOMAS YOUNG, M.D.

CAVAN and CLARE. By HENRY SENIOR.

**CENTRE.** By John Robison, late Professor of Natural Philosophy in the University of Edinburgh.

**CEYLON.** By J. CAPPER, Ceylonese Commissioner to the Great Exhibition of 1851.

CHALMERS, (THOMAS.) By Rev. WILLIAM HANNA, LL.D.

CHEMISTRY. By WILLIAM GREGORY, Professor of Chemistry in the University of Edinburgh.

CHESS. By James Donaldson.

CHILI. By C. B. BLACK.

CHINA. By Sir John Barrow.

CHIVALRY. By Sir Walter Scott, Bart.

CHLOROFORM. By J. Y. SIMPSON, M.D., Professor of Midwifery in the University of Edinburgh.

CHRONOLOGY.

CIVIL LAW. By DAVID IRVING, LL.D.

CLIMATE. By Sir John Leslie.

#### VOLUME VII.

CLOCK AND WATCH WORK. By EDMUND BECKETT DENISON, M.A., Q.C.

COHESION. By Thomas Young, M.D., F.R.S.

COINAGE. By ROBERT MUSHET of the Royal Mint.

COLD and DEW. By Sir John Leslie, late Professor of Natural Philosophy in the University of Edinburgh.

collision, combination, corn-laws, corn-trade, and cottage system. By J. R. M'Culloch.

COLLIERY. By WILLIAM ALEXANDER, Mining Engineer.

COLONY. By James Mill.

COLOUR-BLINDNESS. By George Wilson, M.D., F.R.S.E.

COMET. By Thomas Galloway, F.R.S.

COMMUNISM and CORPORATION. By J. H. BURTON.

CONIC SECTIONS. By WILLIAM WALLACE, LL.D., late Professor of Mathematics in the University of Edinburgh.

CONSTRUCTION. By WILLIAM HOSKING, Author of "Architecture."

**CONSTANTINOPLE.** By EDWARD SANG, late Professor of Mechanical Philosophy in the Imperial School, Constantinople.

COPPER SMELTING. By JAMES NAPIER, Glasgow.

CORK. By HENRY SENIOR.

COTTON MANUFACTURE. Revised by THOMAS BAZLEY, Chairman of the Chamber of Commerce, Manchester.

COWPER. By A. SMITH, Author of "A Life Drama," &c.

CRIMEA. By James Laurie.

CRUSADES, CUVIER, DANTON, and DEFOE. By JAMES BROWNE, LL.D.

CRUSTACEA. By JOHN FLEMING, D.D., Professor of Natural Science, New College, Edinburgh.

CUNNINGHAM, DALRYMPLE, and DEMPSTER. By DAVID IRVING, LL.D.

DAIRY. By John Wilson, Author of "Agriculture."

DEAF and DUMB. By P. M. ROGET, M.D., F.R.S.

**DEMOSTHENES.** By WILLIAM SPALDING, M.A., Professor of Logic and Metaphysics in the University of St Andrews.

**DENMARK.** By EMERIC SZABAD, Author of the Article "Austria."

DIALLING. By HENRY MEIKLE, C.E.

Commerce—Constantinopolitan History—Copyright—Crystallization—Deluge.

## VOLUME VIII.

DIPLOMACY and ENTAIL. By JOHN HILL BURTON.

DISTILLATION. By James Stark, M.D., F.R.S.E.

DIVING and DIVING BELL. By George Buchanan, F.R.S.E.

DOCK and DOCK-YARD. By Sir John Barrow, Bart. Revised by his Son, John Barrow, Admiralty.

**DOLLOND, DOLOMIEU, and DUHAMEL.** By Thomas Young, M.D, F.R.S.

DONEGAL, DOWN, and DUBLIN. By HENRY SENIOR.

**DRAINAGE of TOWNS.** By W. Hosking, Author of "Architecture," &c.

DRAINAGE of LANDS. By J. WILSON, Author of "Agriculture," &c.

DRAMA. By Sir Walter Scott, Bart.

DRAWING and ENGRAVING. By W. H. LIZARS.

DRY ROT. By Sir John Barrow, Bart.

DUMONT, LORD DUNCAN, &c. By James Browne, LL.D.

**DUNBAR**, (WILLIAM,) and ENGLISH LANGUAGE. By DAVID IRVING, LL.D.

DYEING (Calico-Printing.) By F. CRACE CALVERT, Professor of Chemstry, Royal Institution, Manchester.

**DYNAMICS.** By John Robison, late Professor of Natural Philosophy in the University of Edinburgh.

ECONOMISTS. By James Mill.

EDINBURGH and EDINBURGHSHIRE.

EGYPT. By R. STUART POOLE, of the Department of Antiquities, British Museum.

ELECTRICITY. By Sir David Brewster, K.H., M.A., D.C.L, &c.

EMBANKMENT. By J. C. Loudon.

EMIGRATION. By J. R. M'CULLOCH, Author of "Commercial Dictionary," &c.

ENGLAND. By JAMES BROWNE, LL.D., with Continuation.

#### VOLUME IX.

ENTOMOLOGY, FISHERIES, and EDWARD FORBES. By JAMES WILSON, F.R.S.E.

EPHRAEM SYRUS. By Rev. HENRY BURGESS, LL.D.

EPISCOPACY. By Rev. GEORGE GLEIG, D.D.

ERASMUS and FEUDAL LAW. By DAVID IRVING, LL.D.

EQUATIONS. By JAMES IVORY, F.R.S.

ETHNOLOGY. By R. G. LATHAM, M.A., M.D.

ETRUSCANS, EUGENE, FENELON, &c. By James Browne, LL.D.

**EUROPE.** By Charles MacLaren, F.R.S.E. and James Laurie.

EVIL. By Rev. W. L. ALEXANDER, D.D.

EXAMINATIONS. By J. F. MACLENNAN.

EXCHANGE, EXCHEQUER BILLS, and EXCISE. By J. R. M'Culloch.

EXTREME UNCTION, FATHERS, FEDERAL GOVERN-MENT, &c. By Rev. J. TAYLOR, D.D.

FABLE and FALLACY. By WILLIAM SPALDING, A.M., Professor of Logic in the University of St Andrews.

FALCONER, FARQUHAR, and FAIRFAX. By ROBERT CARRUTHERS.

FASHION. By Dr Doran, Author of "Habits and Men," &c.

FERMANAGH. By HENRY SENIOR.

FEZZAN. Revised by Augustus Petermann.

FICHTE. By John Colquioun, F.R.S.E.

FIFESHIRE and FORTH. By THOMAS BARCLAY.

FIGURE of the EARTH. By THOMAS GALLOWAY, F.R.S.

FILTER. By George Buchanan, F.R.S.E.

FLINTSHIRE. By John Girdwood.

FLORIDA. By J. Smith Homans, New York.

FLUXIONS. By WILLIAM WALLACE, LL.D.

FONTANA, FOSTER, and FOURCROY. By Thomas Young, M.D.

FOOD. By Thomas Lindley Kemp, M.D.

FORFAR. By JAMES COWIE.

FORTIFICATION. By Colonel Portlock, Woolwich.

FOSTER. By J. E. RYLAND, M.A.

FOX, C. J. By John Allen.

Exhibition-Fanaticism.

## VOLUME X.

FRANCE. By A. V. KIRWAN, of the Middle Temple, Barrister at Law.

FRANKLIN, (BENJAMIN.) By ALEXANDER NICHOLSON.

FRANKLIN, Sir JOHN. By Sir JOHN RICHARDSON.

FUEL and GAS LIGHT. By CHARLES TOMLINSON, Editor of "Cyclopædia of Useful Arts," &c.

FULLER, ANDREW. By J. E. RYLAND, M.A.

FUNDING SYSTEM. By D. RICARDO, supplemented by J. I. RICARDO, M.P.

FURNACE. By GEORGE BUCHANAN, F.R.S.E.

GALILEO. By JAMES BROWNE, LL.D.

GALWAY. By HENRY SENIOR.

GANGES. By EDWARD THORNTON, India House.

GASSENDI and GIBBON. By HENRY ROGERS, Author of the "Eclipse of Faith," &c.

GEOGRAPHY. By Rev. John Wallace, D.D.

GEOMETRY. By WILLIAM WALLACE, LL.D.

GEOMETRY, ANALYTICAL. By Rev. P. Kelland, M.A., Professor of Mathematics in the University of Edinburgh.

GERMANY. By W. JACOB. Revised by JAMES LAURIE.

GLACIER and FRESNEL. By J. D. FORBES, Professor of Natural Philosophy in the University of Edinburgh.

GLASGOW. By John Strang, LL.D.

GLASS. By James Ballantyne.

GNOSTICISM. By JOHN TULLOCH, D.D., Primarius Professor of Divinity, St Andrews.

GOETHE. By Thomas de Quincey.

GOLDSMITH. By the Right Hon. T. B. MACAULAY.

CAPE of GOOD HOPE. By B. C. PINE, late Lieutenant-Governor of Natal.

GOTAMA BUDDHA. By Rev. R. SPENCE HARDY, Hon. M.R.A.S., Author of "Eastern Monachism," &c.

GOTHS. By LEONARD SCHMITZ, LL.D.

GOVERNMENT. By P. E. Dove, Author of the "History of Human Progression," &c.

GRAMMAR. By Bishop GLEIG. Revised by W. SPALDING, Professor of Logic in the University of St Andrews.

## VOLUME XI.

GRAY, HERRICK, and HOGG. By ROBERT CARRUTHERS.

GREECE. By CHARLES MACLAREN, F.R.S.E. Revised.

GREEK CHURCH. By W. M. HETHERINGTON, D.D., LL.D.

GREGORY of NAZIANZUM. By JOHN TULLOCH, D.D.

GREGORY (Dr. JAMES). By W. P. ALISON, M.D.

GUINEA and HOUSSA. By Augustus Petermann, F.R.G.S., &c.

GUN-COTTON, GUNPOWDER, GUTTA PERCHA, and HAT-MAKING. By CHARLES TOMLINSON.

GUNMAKING. By P. E. Dove.

GUNNERY. By Colonel PORTLOCK.

HALL (ROBERT). By HENRY ROGERS.

HARBOURS. By Thomas Stevenson, C.E.

HARE (C. J.) By W. L. ALEXANDER, D.D.

HARVEY. By Thomas LAYCOCK, M.D., Professor of the Practice of Physic in the University of Edinburgh.

**HEAT.** By T. S. Traill, M.D., Professor of Medical Jurisprudence in the University of Edinburgh.

HELMINTHOLOGY. By JAMES WILSON, F.R.S.E.

HEMP. By T. C. ARCHER, Author of "Popular Economic Botany," &c.

HERALDRY. By T. W. King, York Herald, Herald's College.

HEYNE. By Sir WILLIAM HAMILTON, Bart.

HIEROGLYPHICS. By R. S. Poole, M.R.S.L., &c.

HIMALAYA MOUNTAINS. By Joseph D. Hooker, M.D., F.R.S.

HINDUSTAN. Revised by EDWARD THORNTON, India House.

HISTORY. By DAVID MASSON, M.A., Professor of English
Literature, University College.

HOLLAND. Revised by the Rev. James Ingram, M.A.

HOMER. By JOHN S. BLACKIE, Professor of Greek in the University of Edinburgh.

HOMEOPATHY. By W. T. GAIRDNER, M. D.

HOOD (THOMAS). By RICHARD MONCKTON MILNES.

HORACE. By Theodore Martin.

HORSE, HORSEMANSHIP, and HOUND. Revised by W. H. Langley, Editor of "Bell's Life in London."

HORTICULTURE. By CHARLES MACINTOSH, Author of the "Book of the Garden."

HOWARD (JOHN). By HEPWORTH DIXON.

HOUSEHOLD (ROYAL). By SAMUEL REDGRAVE.

In Two Volumes Medium 8vo, Price £3, with a Complete Index, and upwards of 554 Illustrations,

## A NEW EDITION OF

# KITTO'S CYCLOPÆDIA

OF

# BIBLICAL LITERATURE.

EDITED BY

## THE REV. HENRY BURGESS, LLD.

Compiled by numerous able Scholars and Divines, British, Continental, and American, whose Initials are affixed to their respective Contributions.

The following is a Conspectus of the Work:—

- 1. BIBLICAL CRITICISM—Embracing the History of the Bible Languages—The Canon of Scripture—Literary History and Peculiarities of the Sacred Books—Formation and History of the Texts of Scripture.
- 2. BIBLICAL INTERPRETATION Comprehending the Principles of Translating and Expounding Scripture—Critical Estimate of the leading Interpreters and Principal Versions, Ancient and Modern.
- 3. HISTORY—Proper Names of Persons—Biographical Sketches of the more Prominent Characters—Detailed Accounts of the Principal Events recorded in Scripture—Chronology and Genealogy of Scripture.
- GEOGRAPHY—Names of Places—Description of Scenery— Boundaries and Mutual Relations of the Countries mentioned in Scripture, so far as necessary to illustrate the Sacred Text.
- 5. ARCHÆOLOGY Manners and Customs of the Jews, and other Nations mentioned in Scripture—their Sacred Institutions—their Military Affairs—their Political Arrangements—their Literary and Scientific Pursuits.
- 6. PHYSICAL SCIENCE—Scripture Cosmogony and Astronomy—Zoology—Mineralogy—Botany—Meteorology.

# ATLASES.

# LAST EDITIONS, WITH ALL THE LATEST DISCOVERIES.

GENERAL ATLAS OF THE WORLD. Last Edition, containing all the Latest Discoveries. Maps of the Crimea, the Baltic, Arctic Regions, Australian and African Colonies, on an extended scale; some of the principal of the United States of America; and an Index of 60,000 Names, Imperial Folio, half-bound, price £2:16s.

HAND-ATLAS OF MODERN GEOGRAPHY. A Series of Thirty-Eight Maps. By WILLIAM HUGHES, F.R.G.S.; and JOHN BARTHOLOMEW. With a Complete Index of Names. Quarto, half-bound, price 21s.

SCHOOL ATLAS OF MODERN GEOGRAPHY. New Edition. A Series of Thirty-Seven Maps, for Instruction in Physical, Ancient, and Scripture Geography. With Index. 4to or 8vo, cloth, price 10s. 6d.

"In comprehensiveness, accuracy, finished execution, jud cious adaptation to educational purposes, and moderateness of price, this Atlas stands quite alone."—ATHENÆUM.

"This is, without exception, one of the best Atlases we ever saw. We commend this Atlas strongly."—English Journal of Education.

SCHOOL ATLAS FOR BEGINNERS. New Edition. A Series of Twenty-Seven Maps of the Principal Countries in the World. Oblong 12mo or 18mo, price 2s. 6d.

"Deserves a wide circulation. The Maps by Mr. Bartholomew are admirably executed. Compared with other Atlases of the same size and price which we have seen, this has decidedly the advantage."—Athenæum.

#### v.

ATLAS OF AUSTRALIA, WITH ALL THE GOLD REGIONS. A Series of Maps from the latest and best authorities; beautifully engraved and coloured, and strongly bound in cloth, price 5s.

I. General Map of Australasia, New Zealand, Polynesia and Surrounding Islands. II. Australia.—Divided into Districts. III. New South Wales, Victoria, and South Australia, shewing the different Gold Deposits. IV. New South Wales—Divided into counties, with all the Gold Deposits accurately laid down, and a plan of Sydney. V. Victoria—Divided into Counties, with all the Gold Districts accurately laid down, and a plan of the Mount Alexander Gold Region. VI. New Zealand, Van Diemen's Land, and the settled portion of Western Australia, comprehending Swan River and Australind.

# BLACK'S GUIDE BOOKS

FOR

# ENGLAND, SCOTLAND, AND IRELAND.

Illustrated by Maps, Charts, numerous Views of the Scenery, full particulars regarding Hotels, Inns, Rates of Charges, Distances, and every information likely to prove Useful or Instructive.

## IN PORTABLE VOLUMES, STRONGLY BOUND IN CLOTH.

"We have looked carefully through the volumes. They are admirably 'got up;' the descriptions are accurate, and remarkably clear and comprehensive. We have seldom examined books better 'edited.' . . . . Altogether, this series of works is of immense value to Tourists."—ART JOURNAL. "They should find a corner in the portmanteau of every person about to undertake a journey of pleasure or business either in England and Wales, or Scotland."—JOHN BULL.

- ENGLAND, Fourth Edition, Corrected throughout, and greatly Enlarged, containing 198 Routes, and Illustrated by Twenty-seven elaborately constructed Maps and Charts, besides Views of the Scenery; an Alphabetical List of Places; Exhibitions, &c., in London, with the Mode of obtaining Admission; all the Hotels and Inns, Rates of Charges, and a Comprehensive General Index. Price 10s. 6d.
- SCOTLAND, Twelfth Edition, Greatly Improved, and Illustrated by One hundred and thirty-six Engravings, consisting of an Accurate Travelling Map, Engraved Maps and Charts of Roads, Railroads, and interesting Localities, Plans of the towns of Edinburgh, Glasgow, Perth, and Aberdeen, Views of the Scenery, after Montague Stanley, Cattermole, Leitch, Thomson, J. M. Richardson junior, Sir Thomas Dick Lauder, &c.; and containing full information regarding the best Hotels, Rates of Charges, Coach and Steamer Routes, Distances, and other matters affecting the Convenience of the Tourist. Price 8s. 6d.
- IRELAND, embracing descriptions of all the Routes frequented by Tourists, with full information regarding Hotels, Inns (and the charges made by each), Conveyances, Guides, and every Topographical Information likely to prove useful. Illustrated by a Map of Ireland. Chart of the Lakes of Killarney on a large scale, and Plans of the principal towns. In a neat pocket volume, bound in cloth limp, price 5s.

  The work is divided into four sections, each of which may be had separately at 1s. 6d. each, bound in limp cloth.
- I.—DUBLIN AND THE WICKLOW | III.—THE SHANNON ANDWEST. | IV.—BELFAST, GIANT'S CAUSE-II.—KILLARNEYANDTHE SOUTH. | WAY, AND THE NORTH.

- ENGLISH LAKES, Seventh Edition, Enlarged and Improved. Including an Essay on the Geology of the District, by John Phillips, M.A., F.R.S., F.G.S., Deputy Reader in Geology in the University of Oxford; and Memoranda for Botanists. With a minutely accurate Map, by W. Hughes; Charts of the Lakes, Twelve Outline Views of Mountain Groups, and Views of the Scenery on Wood and Steel. Containing, also, an ample Itinerary of all the Routes, with the Distances accurately laid down, all the best Hotels and Inns, and a Complete Index. Price 5s. (A Cheap Edition of this is published, price 1s.)
- WALES—North and South, and Monmouthshire, Sixth Edition.
  Containing a Full Description of Every Remarkable Place, with Thirtyfive Illustrations, including Maps, Charts, and Views of Scenery; with
  all the Hotels and Inns, and a Copious Itinerary. Price 5s.
- DERBYSHIRE, including Matlock Bath, Chatsworth, Buxton, Castleton, Dovedale, and every other place of interest. With a Map of the County and Plan of Chatsworth. Price 1s.
- HAMPSHIRE AND DORSETSHIRE, including descriptions of the Isle of Wight, Winchester, Southampton, Netley Abbey, Portsmouth, the New Forest, Weymouth, Dorchester, Poole, and every other place of interest, with Map. Price 1s.
- DEVONSHIRE AND CORNWALL, EXETER, PLYMOUTH AND DEVONPORT, LAUNCESTON, TRURO, LANDS-END, AND THE SCILLY ISLANDS. With Map. Price 1s.
- HIGHLANDS AND ISLANDS OF SCOTLAND, including ORKNEY AND ZETLAND. Descriptive of their Scenery, Statistics, Antiquities, and Natural History. Containing, also, Directions for visiting the Lowlands of Scotland, with Maps, Views, Tables of Distances, Notices of Inns, &c. &c. By George and Peter Anderson of Inverness. Third Edition, carefully Revised, Enlarged, and Remodelled. Price 10s. 6d.
- THE TROSACHS, LOCH KATRINE, LOCH LOMOND, and Neighbouring Routes, including the Districts of Aberfoyle, Balquhidder, Menteith, and Rob Roy's Country, &c. &c.; with Map and Numerous Illustrations. Price 1s.
- HIGHLANDS OF PERTHSHIRE, including descriptions of Perth and Environs—Dunkeld, Blair-Atholl, Kenmore, Loch Tay, Killin, Lochearnhead, Callander and the Trosachs; with Map, Charts, and numerous Illustrations. Price 1s.
- HIGHLANDS OF ARGYLESHIRE AND BUTE. Containing a description of Glasgow, Gareloch, Loch Long, Loch Goil, Bute and Arran, and all the principal places of interest and resort on the West Coast of Scotland, the various Steamer and Coach Routes, &c.; with Map. Price 1s.
- STAFFA, IONA, GLENCOE, AND CALEDONIAN CANAL, including a description of Oban and its vicinity; with Map of Mull, and a view of Fingal's Cave. Price 1s.

- ISLAND OF SKYE, and West of Ross-shire; with a Map of Skye. Price 1s.
- ABERDEEN, BALMORAL, AND BRAEMAR, including Excursions among the Cairngorm Mountains; with Map and Illustrations. Price 1s.
- EDINBURGH, with a Description of the Environs, including Melrose, Abbotsford, Dryburgh, St. Andrews, Linlithgow, Lochleven, &c. New Edition. Illustrated with a Plan of the City, a Map of the Country Ten Miles round, and numerous Views of the Public Buildings and Scenery. Price 2s. 6d., or without Illustrations, 1s.
- GLASGOW and the West Coast. Including the Falls of the Clyde, Bute, Arran. Staffa. Iona, and the Land of Burns; with a Plan of Glasgow and other Charts, numerous Views of the Public Buildings and neighbouring Scenery. Price 2s. 6d., or without Illustrations, 1s.
- MOFFAT AND VICINITY, St. Mary's Loch, the Grey Mare's Tail, Loch Skene, &c., and Hints to Anglers in the Rivers, Streams, and Lochs in the Neighbourhood; with Map and Illustrations. Price 1s.

## BLACK'S ROAD AND RAILWAY TRAVELLING MAPS.

- Carefully constructed from the Maps of the Ordnance Survey and other Authorities, and containing all the Roads, Railroads, Villages, Country Seats, Fishing Streams, Rivers, Lakes, and Mountains, and every Topographical information required by the Tourist on pleasure or business. Lined with, or printed on, patent cloth, and neatly bound in portable cases.
- ENGLAND AND WALES. 32 Inches by  $22\frac{1}{2}$ . 4s. 6d. Do. Smaller. Size 19 Inches by 15, 2s. 6d. Or uncoloured and unmounted, 1s.
- ENGLISH LAKE DISTRICT. 19 Inches by 14. 2s. 6d. Do. Do. Uncoloured and unmounted. Price 8d.
- WALES (North and South). 14 Inches by  $11\frac{1}{4}$ . 1s. 6d. each.
- SCOTLAND. 32 Inches by 22½. Price 4s. 6d. Do. Smaller. Size 19 Inches by 15, 2s. 6d. Or uncoloured and unmounted, 1s.
- IRELAND. Size, 20 Inches by  $14\frac{1}{2}$ . Price 2s. 6d.
- TOURIST'S AND SPORTSMAN'S COMPANION TO THE COUNTIES OF SCOTLAND. A Series of Thirty-six Maps, shewing all the Roads, Railways, Villages, Country Seats, Rivers and Lakes, Places of Historical and Legendary Note; Memoranda of Battles, Heights of Mountains, &c. &c. In a portable volume, strongly bound. Price 10s. 6d.
- MEMENTO OF THE TROSACHS, LOCH KATRINE, AND LOCH LOMOND. A Series of Twenty-seven Engravings, by BIRKET FOSTER. Square 18mo. Price 1s., and in Cloth, 1s. 6d.

# RELIGIOUS WORKS.

- THE GOSPEL IN EZEKIEL: Illustrated in a Series of Discourses. By Thomas Guthrie, D.D., Author of "Pleas for Ragged Schools," &c. Crown 8vo, price 7s. 6d. Twelfth Thousand.
  - "He dips his brush in the simplest colours, and as with a firm hand, bold and strong, he fills his canvas, figures start into life there that every eye at once recognises; and in attitudes whose meaning needs no interpreter to describe. Their author has looked on the material world with the eye of a poet, on human life with the eye of a philanthropist, the two great open fountains whence his imagery as a pulpit orator is drawn. A volume that will earn attention for its sacred themes, wherever taste for pictorial writing exists, which our current literature has created; and which is destined, we believe, by reason of its more enduring merits, to pass into the hands of our children's children, as one of the manuals of piety by which, in future generations, the faith of our Scottish people will be brightened and refreshed."—NORTH BRITISH REVIEW.
  - "To our friends south of the Scottish Border, who do not know Dr. Guthrie, we say, procure this volume, and read it, and you will feel that you have made the acquaintance of a man whom it were worth while to go some distance to see."—British Quarterly REVIEW.

- BERRIDGE'S CHRISTIAN WORLD UNMASKED. New Edition. Edited, and with a Life of the Author. By Thomas GUTHRIE, D.D. Foolscap 8vo, price 2s. 6d.
  - "The Christian World Unmasked has been long known and highly esteemed. Its racy style, bold imagery, profound analysis of the human heart unrenewed by the grace of God, and faithful application of the truth to the conscience, make it one of the best books ever written, and one of the most profitable any one can read. The present edition is enriched by 'A Life of the Author,' from the pen of one who claims kindred genius and eloquence with John Berridge."—WITNESS.

- THE HISTORY OF PALESTINE: From the Patriarchal Age to the present time. By John Kitto, D.D., Author of Popular Dictionary of the Bible, &c. Profusely Illustrated. Crown 8vo, cloth plain, 6s.; or cloth, gilt edges, 6s. 6d.
  - "Beyond all dispute it is the best historical compendium of the Holy Land, from the days of Abraham to those of the late Pasha of Egypt, Mehemet Ali."—OBSERVER.

" Not only a complete description of the Holy Land, but a condensed History of the Jewish People."-JOHN BULL.

#### Religious Works-Continued.

#### IV.

A TREATISE ON BIBLICAL CRITICISM. Exhibiting a Systematic View of that Science. By Samuel Davidson, D.D., Author of "Ecclesiastical Polity," &c. New Edition. In One Volume 8vo, price 18s.

"The present work bears marks of the same industry and learning which have distinguished the author's other productions."—BIB. SACRA.

"We have no hesitation in pronouncing this work to be the most important contribution ever made in this country to the science of Biblical Criticism."—CLERICAL JOURNAL.

#### V.

A POPULAR DICTIONARY OF THE BIBLE. By John Kitto, D.D., Author of the "Cyclopædia of Biblical Literature," &c. In One Volume 8vo, illustrated by 336 Engravings on Wood, price 10s. 6d.

This work is studiously accommodated to the wants of the great body of the religious public. To Parents, to Sunday School Teachers, to Missionaries, and to all engaged, either statedly or occasionally, in the important business of Biblical education, the volume is confidently recommended as the most valuable compendium of Bible knowledge for the people which has ever appeared in this country.

#### VI.

DR. WARDLAW'S LIFE AND CORRESPONDENCE. By W. L. ALEXANDER, D.D. Second Edition. In One Volume, demy 8vo, Price 12s.

"We recommend this work in the most earnest and unqualified manner, persuaded as we are that it will take its permanent rank among the comparatively few of those biographical works which we not only read again and again, and place on the shelf of favourite authors, but also bequeath and recommend to children."—EVANGELICAL MAGAZINE.

"The book sufficiently recommends itself. It is worthy of the good and great name which it embalms, and both from that name and its own masterly execution, will speedily attain a wide circulation, be read with eager interest, and then laid up in store. Successive generations of Christians will preserve it with their choicest treasures and legacies from the present."—Eclectic Review.

#### VII.

RALPH WARDLAW'S SYSTEMATIC THEOLOGY. Edited by the Rev. J. R. CAMPBELL, M.A. In 8vo, Price 12s., Vol. I. (To be completed in 3 Vols).

# SIR WALTER SCOTT'S WRITINGS AND LIFE.

WAVERLEY NOVELS in Sets. Five Editions as follows:—

- I. LIBRARY EDITION, uniform with the Standard English Authors. Illustrated by upwards of Two Hundred Engravings on Steel, after Drawings by Turner, Landseer, Stanfield, Wilkie, Roberts, Frith, Ward, Philips, Elmore, Faed, &c. In 25 vols. demy 8vo, elegantly bound in extra cloth, gilt, price £13:2:6.
- II. ABBOTSFORD ILLUSTRATED EDITION. With
   120 Engravings on Steel, and nearly 2000 on Wood.
   12 vols. super-royal octavo, cloth, £14:14s.
- III. AUTHOR'S FAVOURITE EDITION in 48 small-sized vols. With 96 Engravings, £7:4s.
- IV. CABINET EDITION. In 25 small-sized vols., £3, 13s. 6d.
- V. PEOPLE'S EDITION. 5 large vols. royal 8vo, cloth, £2:2s., or each novel separate, sewed, price 1s. 6d.

SIR WALTER SCOTT'S WRITINGS AND LIFE—Continued.

POETICAL WORKS. Six Editions as follows:—

- I. In one small volume. With Illustrations on Wood and Steel. Gilt edges, 5s.; or on larger paper, with additional Engravings, 6s.
- II. POCKET EDITION FOR TOURISTS. LAY OF THE LAST MINSTREL—MARMION—LADY OF THE LAKE—ROKEBY—and LORD OF THE ISLES. 1s. 6d. each; or bound in morocco, 2s. 6d.
- III. In one large vol. royal 8vo, double cols. 10s.; or large paper, with 26 Engravings from Turner, 18s.; full morocco, elegant, 32s.
- IV. In Six small vols. 12 Engravings after Turner, 24s.
- V. In Twelve small vols. 24 Illustrations by Turner, £1:16s.
- VI. ILLUSTRATED EDITIONS of THE LADY OF THE LAKE, LAY OF THE LAST MINSTREL, MARMION, and THE LORD OF THE ISLES. Exquisitely illustrated by Birket Foster and John Gilbert. Extra cloth, gilt edges, 18s.; Morocco, elegant or antique, 25s.

# PROSE WORKS. Two sets as follows:—

- I. In 28 vols. 56 Engravings from Turner, &c., £4:4s.
- II. In 3 vols. royal 8vo, uniform with the People's Edition, £1:6s.

# SIR WALTER SCOTT'S WRITINGS AND LIFE—Continued. TALES OF A GRANDFATHER. (History of Scotland).

- I. In 3 small vols., illustrated, cloth, 12s.; extra, gilt edges, 15s.
- II. In one large vol. royal 8vo, double cols., 6s.; or large paper, with 11 Engravings after Turner, 10s. 6d.
- III. School Edition. 2 small vols. 10s.
- IV. HISTORY OF FRANCE. One small volume, 4s.; gilt edges, 5s.

## LIFE OF NAPOLEON BONAPARTE.

I. In 5 small vols. with Maps, Portraits, and 9 Engravings after Turner, 20s. II. In one large vol. royal 8vo, double cols., 10s.; or large paper, with 23 Engravings after Turner and others, 18s.

# LIFE OF SIR WALTER SCOTT. By J.

G. LOCKHART, Esq. I. In one small vol. 12 Engravings, 7s. 6d.; extra gilt edges, 8s. 6d. II. In one large vol. royal 8vo, double cols., 10s.; or with 11 Engravings from Turner and others, 18s. III. In Ten small vols. 20 Engravings, 30s.

# BEAUTIES OF SIR WALTER SCOTT—

A Selection from his Writings, 5s.; extra, gilt edges, 6s.

READINGS FOR THE YOUNG, from the Works of Sir Walter Scott. Numerous Illustrations, 7s.; or in 3 separate vols., 2s. 6d. each.

# WORKS ON GARDENING AND BOTANY.

I.

THE COTTAGE GARDEN. By Robert Adamson. 12mo, price 2s. Second Edition.

II:

NEILL'S FRUIT, FLOWER, AND KITCHEN GARDEN. Fifth Edition, improved, with Additions, price 5s.

"In Horticultural Literature, no book has acquired a higher reputation than the late Dr. Neill's concise and popular treatise on Practical Gardening. It is copiously illustrated; and no gardener, be he amateur or professional, great or small, whether he grows pines or only potatoes, should be without it."—MORNING POST.

III.

THE AMATEUR GARDENER'S YEAR-BOOK: A Guide for those who cultivate their own Gardens in the Principles and Practice of Horticulture. By the Rev. Henry Burgess, LL.D. and Ph. D., &c. &c. Fcap. 8vo, price 5s., with illustrated Frontispiece and Title.

"This is, beyond compare, the best work of its class that has come under our notice. It is really popular."—CRITIC

under our notice. It is really popular."—CRITIC.
"Contains a great deal of valuable information in laying out gardens, and keeping a suitable succession of flowering plants and shrubs."
—ATLAS.

IV.

OUTLINES OF BOTANY: Being an Introduction to the Study of the Structure, Functions, Classification, and Distribution of Plants. With a Glossary and Copious Index. By John Hutton Balfour, M.D., F.R.S.L. and E., F.L.S., Professor of Botany in the University of Edinburgh. Fcap. 8vo, 7s. 6d., illustrated by 595 Wood Engravings.

٧.

CLASS-BOOK OF BOTANY: Being an Introduction to the Study of the Vegetable Kingdom. By J. H. BALFOUR, M.D., F.R.S.L. and E., Regius Keeper of the Royal Botanic Garden, Professor of Medicine and Botany in the University of Edinburgh. In one large Volume 8vo, with 1800 Illustrations, price 31s. 6d.

The same may also be had in Two Parts.

- PART I.—STRUCTURAL & MORPHOLOGICAL BOTANY, 8vo, 10s. 6d.
- PART II.—COMPRISING THE ELEMENTS OF VEGE-TABLE PHYSIOLOGY, CLASSIFICATION, BOTANI-CAL GEOGRAPHY, AND FOSSIL BOTANY, WITH A GLOSSARY OF TERMS. 8vo. 21s.

# MEDICAL AND SURGICAL WORKS.

I.

By JAMES MILLER, F.R.S.E.,

Professor of Surgery in the University of Edinburgh, &c. &c. THE PRINCIPLES OF SURGERY. Third Edition, illustrated by 240 Engravings. 8vo. 16s.

II.

By the same Author,

THE PRACTICE OF SURGERY. Third Edition, illustrated by 227 Engravings. 8vo. 16s.

III.

By Sir George Ballingall,

Late Professor of Military Surgery in the University of Edinburgh, OUTLINES OF MILITARY SURGERY. Fifth Edition, illustrated. 8vo. 14s.

IV.

By WILLIAM GREGORY,

Professor of Chemistry in the University of Edinburgh, ELEMENTARY TREATISE ON CHEMISTRY. Illustrated. Fcap. 8vo. 5s.

v.

By Robert Christison, M.D.,

Professor of Materia Medica in the University of Edinburgh, A TREATISE ON POISONS. In relation to Medical Jurisprudence, Physiology, and the Practice of Physic. Fourth Edition, enlarged and improved. 8vo. 10s.

VI.

By John Hutton Balfour, M.D.,

Professor of Medicine and Botany in the University of Edinburgh, CLASS-BOOK OF BOTANY. Being an Introduction to the Study of the Vegetable Kingdom. 8vo. 31s. 6d.

VII.

By Thomas Laycock, M.D.,

Professor of the Practice of Medicine in the University of Edinburgh, &c.,

LECTURES ON THE PRINCIPLES AND METHODS OF MEDICAL OBSERVATION AND RESEARCH; for the use of Advanced Students and Junior Practitioners. Crown 8vo, price 6s.

## MEDICAL AND SURGICAL WORKS-Continued.

#### VIII.

## By ALEXANDER MACAULAY, M.D.,

A DICTIONARY OF MEDICINE, designed for Popular Use. Containing an Account of Diseases and their Treatment, including those more frequent in Warm Climates, with directions for administering the various substances used as Medicines, the Regulation of Regimen and Diet, and the management of the Diseases of Women and Children. Thirteenth Edition. 8vo. 12s.

#### IX.

By James Y. Simpson, M.D., F.R.S.E.,
Professor of Medicine and Midwifery in the University of
Edinburgh,

OBSTETRIC MEMOIRS AND CONTRIBUTIONS, including those on Anæsthesia. Edited by W. O. PRIESTLEY, M.D., and H. R. STORER, M.D. 2 vols. 8vo. 36s.

#### X.

## By the same Author,

PHYSICIANS AND PHYSIC. Three Addresses: I. ON THE DUTIES OF YOUNG PHYSICIANS. II. ON THE PROSPECTS OF YOUNG PHYSICIANS. III. ON THE MODERN ADVANCEMENT OF PHYSIC. Crown 8vo. 3s. 6d.

#### XI.

## By T. S. TRAILL, M.D.,

Professor of Medical Jurisprudence in the University of Edinburgh, MEDICAL JURISPRUDENCE. Being Outlines of a Course of Lectures. 12mo. 5s.

#### XII.

## By Charles Wilson, M.D.,

THE PATHOLOGY OF DRUNKENNESS. A View of the Operation of Ardent Spirits in the Production of Disease, founded on Original Observation and Research. Fcap. 8vo. 3s.

#### XIII.

ADDRESSES TO MEDICAL STUDENTS: Delivered at the instance of the Edinburgh Medical Missionary Society, 1855-6. By W. P. Alison, M.D., Emeritus Professor of the Practice of Physic in the University of Edinburgh; George Wilson, M.D., Professor of Technology in the University of Edinburgh; Andrew Wood, M.D., President of the Royal College of Surgeons of Edinburgh; Benjamin Bell, F.R.C.S.E.; and John Coldstream, M.D. Fcap. 8vo. 3s. 6d.

## SCHOOL BOOKS.

- CARSON—EXERCISES IN ATTIC GREEK, for the Use of Schools and Colleges. By A. R. Carson, LL.D., F.R.S.E., &c., and late Rector of the High School of Edinburgh. 12mo, 4s. bound.
- CARSON PHÆDRI FABULÆ, AUGUSTI LIBERTI, FABULARUM ÆSOPIARUM, quas oculis puerorum subjici fas est, libras quinque, cum indice verborum, phrasiumque difficiliorum Anglice redditorum. Edidit. A. R. CARSON, LL.D., late Rector of the High School, Edinburgh. Editio Sexta. 18mo, 2s. bound.
- KELLAND—ELEMENTS OF ALGEBRA. By P. Kelland. A.M., F.R.SS.L. & E., &c. late Fellow of the Queen's College, Cambridge, Professor of Mathematics in the University of Edinburgh. 8vo, 9s. cloth.
- KITTO—HISTORY OF PALESTINE FOR SCHOOLS. From the Patriarchal Age to the present time; with Introductory Chapters on the Geography and Natural History of the Country, and on the Customs and Institutions of the Hebrews. By John Kitto, D.D., F.S.A., Editor of the "Pictorial Bible," &c. With questions for examination, by Alexander Reid, LL.D., Rector of the Edinburgh Institution. 12mo, 3s. 6d., or with Map of Palestine, 4s. bound.
- PILLANS—FIRST STEPS IN THE PHYSICAL AND CLAS-SICAL GEOGRAPHY OF THE ANCIENT WORLD. By James Pillans, Professor of Humanity in the University of Edinburgh. Second Edition, With Maps. 12mo. Price 1s. 6d. cloth.
- PORTEOUS—A SUMMARY OF THE PRINCIPAL EVIDENCES FOR THE TRUTH AND DIVINE ORIGIN OF THE CHRISTIAN REVELATION. By the late Beilby Porteous, D.D. New Edition by Dr. Boyd, late of the High School, Edinburgh. 18mo, 1s. bound.

- SCHMITZ—ELEMENTARY GRAMMAR OF THE GREEK LANGUAGE. By Dr. L. SCHMITZ, F.R.S.E., Rector of the High School, &c. &c. 12mo, 3s. 6d. cloth.
- SCHMITZ—HANDBOOK OF ANCIENT HISTORY, from the earliest times to the overthrow of the Western Empire, (comprising the History of China, India, Bactria, Media, Persia, Assyria, Babylonia, Lydia, Phoenicia, Egypt, Carthage, Greece, Rome, &c. By Dr. Leonhard Schmitz, F.R.S.E., Rector of the High School of Edinburgh. 7s. 6d., or in two parts at 4s. each.
- SCRYMGEOUR—A CLASS BOOK OF ENGLISH POETRY. Comprising Extracts from the most distinguished Poets of this Country, from Chaucer to the Present Time, with Biographical Notices, Explanatory Notes, and an Introductory Essay on the Origin and Progress of the English Language. By Daniel Scrymgeour, of Circus Place School, Edinburgh. 12mo, 4s. 6d. bound; or in Two Parts, price 2s. 6d. each.—Part I. containing the Poets from Chaucer to Otway; Part II. from Prior to Tennyson.
- VEITCH—GREEK VERBS, IRREGULAR AND DEFECTIVE; their Forms, Meaning, and Quantity; embracing all the tenses used by the Greek writers, with references to the passages in which they are found. By Rev. W. Veitch, 12mo, 6s. cloth.
- TYTLER—HISTORY OF SCOTLAND. By PATRICK FRASER TYTLER, Esq. Enlarged and Continued to the Present Time, by the Rev. James Taylor, D.D., and adapted to the Purposes of Tuition by ALEXANDER REID, A.M., LL.D., Rector of the Edinburgh Institution. 12mo, 3s. 6d. bound.
- GENERAL MODERN HISTORY. By ALEXANDER FRASER TYTLER, Lord Woodhouselee, late Professor of Universal History in the University of Edinburgh, with a Chronological Table, thoroughly revised and brought down to the end of the Russian War (1856). 12mo, 3s. bound.
- GENERAL ANCIENT HISTORY. By ALEXANDER FRASER TYTLER, Lord Woodhouselee. With a Chronological Table and Map. 12mo, 3s. bound.

# MISCELLANEOUS WORKS.

- MEMORIALS OF HIS TIME. By HENRY COCKBURN, late one of the Senators of the College of Justice. 8vo, price 14s.
  - "The book of the half-year, both in matter and style." Illustrated London News.
  - "We have seen of late no more useful work for fixing our landmarks in one direction, shewing what grievances it was weakness to retain, and what remedial measures it was worse folly to resist."—TIMES.
  - "Edinburgh has sent out few books so full of entertainment, or so high in value as these Memorials."—ATHENÆUM.
- LIFE OF LORD JEFFREY, late one of the Judges of the Court of Session. By Henry Cockburn, late one of the Senators of the College of Justice. Second Edition. 2 vols. 8vo, 25s.
- THE POOR LAW MANUAL FOR SCOTLAND. By ALEXANDER M'NEEL CAIRD, Esq. Sixth Edition. 7s. 6d.
- CLOCK AND WATCH WORK, from the Eighth Edition of the Encyclopædia Britannica. With an Appendix on the Dipleidoscope, and all the latest improvements. With Numerous Illustrations. Fep. 2s.
- SHIPBUILDING. A Treatise (from the Encyclopædia Britannica) on the Theory and Practice of Naval Architecture. By Augustine F. B. Creuze. 4to, 12s.
- THE BREEDING AND ECONOMY OF LIVE STOCK.

  The result of Forty Years' Practical Experience in the
  Management and Disposal of Cattle, Horses, Sheep; and
  Pigs. By James Dickson. 12mo, 3s. 6d.
- THE GENESIS OF THE EARTH AND OF MAN: A Critical Examination of Passages in the Hebrew and Greek Scriptures; chiefly with a View to the Solution of the Question—Whether the Varieties of the Human Species be of more than One Origin. Edited by REGINALD STUART POOLE, M.R.S.L., &c. Crown 8vo, 5s.

## MISCELLANEOUS WORKS-Continued.

- NORWAY AND ITS GLACIERS, visited in 1851. Followed by Journals of Excursions in the High Alps of Dauphiné, Berne, and Savoy. By James D. Forbes, D.C.L., F.R.S., Sec. R.S., Edin., Corresponding Member of the Institute of France, and Professor of Natural Philosophy in the University of Edinburgh. With Two Maps, Ten Lithographic Views printed in colours by Day and Son, and Twenty-two Wood Engravings. Royal 8vo, 21s.
  - "This is one of those books which we need not blush to present to foreign Philosophers and men of learning as a specimen of the literature of science in England."—Examiner.

## By the same Author.

- THE TOUR OF MONT BLANC AND OF MONTE ROSA.

  Being a Personal Narrative, abridged from the Author's

  "Travels in the Alps of Savoy," &c. Illustrated, 12mo, 5s.
  - "An admirable edition, and the Map of the Mer de Glace most correct and valuable."—Albert Smith.
- TURKEY, ANCIENT AND MODERN. Being an Historical Survey of the Ottoman Empire, from its establishment in Europe to the Present Day. By ROBERT W. FRASER, M.A. Crown 8vo, 7s. 6d.
- MUSICAL COMPOSITION: Its Theory and Practice. With numerous Engravings and copious Musical Illustrations interspersed with the text. 4to, 9s.
  - "A masterly and comprehensive Essay."—ATHENÆUM.
- TREATISES ON SUBJECTS CONNECTED WITH ECONO-MICAL POLICY, with Biographical Sketches of Quesnay, Smith, and Ricardo. By J. R. M'Culloch, Esq. 8vo, 14s.
- MANUAL OF MINERALOGY; OR, THE NATURAL HISTORY OF THE MINERAL KINGDOM. Containing a General Introduction to the Science, and descriptions of the several Species, including the more Recent Discoveries and Chemical Analysis. By James Nicol, F.G.S., Professor of Natural History in Marischal College, Aberdeen. Crown 8vo, cloth, price 6s.


## MISCELLANEOUS WORKS-Continued.

- THE GENTLE SHEPHERD. By ALLAN RAMSAY. New Edition, with numerous Illustrations after David Allan. 18mo, 2s. 6d.
- A TREATISE ON THE STEAM-ENGINE. By John Scott Russell, M.A., F.R.S.E. Illustrated. Post 8vo, 5s.

## By the same Author,

- STEAM AND STEAM NAVIGATION. A Treatise on Nature, Properties, and Applications of Steam and on Steam Navigation. Illustrated. Post 8vo, 9s.
- RUSSIAN LIFE IN THE INTERIOR; OR, THE EXPERIENCES OF A SPORTSMAN. By Ivan Tourghenief of Moscow. Edited by J. D. Meiklejohn. Foolscap, 6s.
- "Abounding in pleasant varieties and sketches from life."—ATHENÆUM.
- THE POETRY AND POETS OF BRITAIN. From Chaucer to Tennyson, with Biographical Sketches, and a rapid View of the Characteristic Attributes of each. Preceded by an Introductory Essay on the Origin and Progress of English Poetical Literature. By Daniel Scrymgeour. Post 8vo, 6s. cloth; 6s. 6d. cloth, gilt edges.
- THE WEALTH OF NATIONS. An Inquiry into the Nature and Causes of the Wealth of Nations. By Adam Smith, LL.D. Edited, and with Life of the Author, by J. R. M'Culloch, Esq. Fourth Edition, corrected throughout, and greatly enlarged. 8vo, 16s.
- THE HISTORY OF SCOTLAND. By PATRICK FRASER TYTLER, Esq. Third Edition. 7 vols. 8vo, £2:12:6.
- THE ROD AND THE GUN. By James Wilson, Esq., F.R.S.E. Being Two Treatises on Angling and Shooting. The latter by the Author of "The Oakleigh Shooting Code." Second Edition, with numerous Engravings on Wood and Steel. Post 8vo, 10s. 6d.





BOUND BY A JOHN CRAY

